

香港考試局
HONG KONG EXAMINATIONS AUTHORITY

一九八二年香港中學會考
HONG KONG CERTIFICATE OF EDUCATION EXAMINATION, 1982

MATHEMATICS (SYLL 1)


MARKING SCHEME

This is a restricted document.

It is meant for use by markers of this paper for marking purposes only.

Reproduction in any form is strictly prohibited.

© 香港考試局 保留版權
Hong Kong Examinations Authority
All Rights Reserved 1982

SOLUTION STEPS	MARKS	NOTES
(5 marks)		ALTERNATIVELY,
$\frac{2+i}{1-3i}$		Let $a+bi = \frac{2+i}{1-3i}$
$= \frac{(2+i)(1+3i)}{(1-3i)(1+3i)}$	2M	$(a+bi)(1-3i) = 2+i$
$= \frac{-1+7i}{10}$	1A	$(a+3b)+(b-3a)i = 2+i$ 1A
$= -\frac{1}{10} + \frac{7}{10}i$	1A	$a+3b = 2$ 1M
	1A	$b-3a = 1$ 1M
	1A	$a = -\frac{1}{10}$ 1A
		$b = \frac{7}{10}$ 1A
(5 marks)		ALTERNATIVELY,
$4^{x-y} = 4$		$\frac{4^{x+y}}{4^{x-y}} = 4$ 1M
$4^{x+y} = 16$		
$x-y = 1$	1A	$4^{2y} = 4$ 1A
$x+y = 2$	1A	$2y = 1$ 1A
Solving,	1M	$y = \frac{1}{2}$ 1A
$x = 1\frac{1}{2}$	1A	$x = 1\frac{1}{2}$ 1A
$y = \frac{1}{2}$	1A	
(5 marks)		
$2x^2 - x < 36$	1A	
$2x^2 - x - 36 < 0$	2A	For factorization
$(2x-9)(x+4) < 0$		
$-4 < x < 4\frac{1}{2}$	2A	Accept $\begin{cases} -4 < x \\ x < 4\frac{1}{2} \end{cases}$ "-4 < x and x < 4 1/2"
(6 marks)		
		$\angle C = 90^\circ$
	2M	$\tan \angle BOC = \frac{2\sqrt{3}}{6} = \frac{\sqrt{3}}{3}$
	1A	$\angle BOC = 30^\circ$ or $\frac{\pi}{6}$
Area of sector	1M+1A	$= \pi(6)^2 \times \frac{30}{360}$ or $\frac{1}{2}(6)^2 \frac{\pi}{6}$
	1A	$= 3\pi$

5. (6 marks)

$$2\sin^2\theta + 5\sin\theta - 3 = 0$$

$$(2\sin\theta - 1)(\sin\theta + 3) = 0$$

$$\sin\theta = \frac{1}{2} \quad \text{or} \quad \sin\theta = -3$$

Rejecting $\sin\theta = -3$,

$$\theta = 30^\circ \quad \text{or} \quad 150^\circ$$

1M+1A
1A
1A
1A+1A

1M for attempting to factorize
For $\sin\theta = \frac{1}{2}$
If a cand. writes $\sin\theta = -3$ only, award 2 marks.
Accept $\theta = 30^\circ, 150^\circ$ or $\theta = 30^\circ$ and 150° .
General solution, no mark

If more than 2 answers given, deduct 1 mark for each wrong answer from the marks obtained in the answer only.

6. (6 marks)

(a) (1, 3), (3, 1), (2, 2)

$$\text{Probability} = \frac{3}{36}$$

$$= \frac{1}{12}$$

1A
1A

For numerator
For denominator

(b) (1, 1), (1, 2), (2, 1)

$$\text{Probability} = \frac{3}{36}$$

$$= \frac{1}{12}$$

1A
1A

For numerator
For denominator

(c) Probability = $1 - \frac{1}{12} - \frac{1}{12}$

$$= \frac{5}{6}$$

1M
1A

ALTERNATIVELY,
(1, 4), (1, 5), (1, 6), ... , (6, 6)

$$\text{Probability} = \frac{30}{36}$$

$$= \frac{5}{6}$$

1A
1A

For numerator
For denominator

If answer not simplified, deduct 1 mark for the whole question.

7. (6 marks)

(a) $x = 360 \times \frac{2}{12}$

$$= 60$$

$$y = 210$$

$$z = 90$$

1M
1A
1A
1A

ALTERNATIVELY,
put $x = 2k$
 $y = 7k, z = 3k$
 $2k + 7k + 3k = 360$

(b) Total number = $240 \times \frac{12}{2}$

$$= 1440$$

1M
1A

ALTERNATIVELY,
No. in Kowloon = 840
No. in N.T. = 360
Total no. = $840 + 360 + 240$
= 1440

8. (a) (10 marks)

$$AC^2 = x^2 + x^2$$

$$= 2x^2$$

$$AB^2 = AC^2 + BC^2$$

$$= 2x^2 + y^2$$

$$2x^2 + y^2 = 9^2$$

$$8x + 4y + 9 = 69$$

i.e. $2x + y = 15$

1M
1M
1A
1A

For Pythagoras' Theorem
For Pythagoras' Theorem

Sub. $y = 15 - 2x$ in $2x^2 + y^2 = 9^2$,

$$2x^2 + (15 - 2x)^2 = 81$$

$$6x^2 - 60x + 144 = 0$$

$$x^2 - 10x + 24 = 0$$

$$(x - 4)(x - 6) = 0$$

1M
1A
1A
1A

For solving Sim. Eqs.

$$x = 4 \quad \text{or} \quad 6$$

$$x = 4, \quad y = 7$$

$$x = 6, \quad y = 3$$

1A
1A

For both values
For both values

ALTERNATIVELY,

Sub. $x = \frac{15 - y}{2}$ in $2x^2 + y^2 = 9^2$,

$$2\left[\frac{15 - y}{2}\right]^2 + y^2 = 9^2$$

$$y^2 - 10y + 21 = 0$$

$$(y - 3)(y - 7) = 0$$

$$y = 3 \quad \text{or} \quad 7$$

$$y = 7, \quad x = 4$$

$$y = 3, \quad x = 6$$

1M
1A
1A
1A
1A
1A

For both values
For both values

(b) (2 marks)

$$\cos\theta = \frac{BC}{AB} \quad \text{or} \quad \tan\theta = \frac{AC}{BC} \quad \text{or} \quad \sin\theta = \frac{AC}{AB}$$

$$\cos\theta = \frac{7}{9} \quad \text{or} \quad \tan\theta = \frac{\sqrt{32}}{7} \quad \text{or} \quad \sin\theta = \frac{\sqrt{32}}{9}$$

$$\theta \approx 39^\circ$$

1M
1A

For both values

11. (a) (4 marks)

Let α, β be the roots of $x^2 - 10x + k = 0$

(i) $\alpha + \beta = 10$ _____ 1A
 $OA + OB = 10$ _____ 1A

(ii) $\alpha\beta = k$ _____ 1A
 $OA \times OB = k$ _____ 1A

(b) (4 marks)

(i) $OM + ON = \frac{1}{2}(OA + OB)$ _____ 1M
 $= 5$ _____ 1A

(ii) $OM \times ON = \frac{1}{2}OA \times \frac{1}{2}OB$ _____ 1M
 $= \frac{k}{4}$ _____ 1A

(c) (4 marks)

(i) From (b), $-p = OM + ON$ _____ 1M
 $p = -5$ _____

$r = OM \times ON$ _____
 $= \frac{k}{4}$ _____ 1M

(ii) $OM = 2,$
 $ON = 3$

$\frac{k}{4} = (2)(3)$ _____ 1M

$k = 24$ _____ 1A

ALTERNATIVELY,

$OM = 2$
 $M = (2, 0)$

Sub. in $y = x^2 - 5x + \frac{k}{4}$ _____ 1M

$0 = 4 - 10 + \frac{k}{4}$

$k = 24$ _____ 1A

12. (a) (7 marks)

$Y = k_1x$ or $Z = k_2x^2$ _____ 1M

$P = Y + Z$ _____ 1M
 $= k_1x + k_2x^2$ _____

$80\ 000 = 20k_1 + 20^2k_2$ _____ 1A

$87\ 500 = 35k_1 + 35^2k_2$ _____

Solving, _____ 1M

$k_1 = 6000$ _____ 1A

$k_2 = -100$ _____ 1A

$P = 6000x - 100x^2$

When $x = 15,$
 $P = 6000(15) - 100(15)^2$
 $= 67\ 500$ _____ 1A

(b) (3 marks)

$P = 6000x - 100x^2$
 $= -100(x^2 - 60x)$
 $= -100[x^2 - 60x + 30^2 - 30^2]$ _____ 2M
 $= -100[(x - 30)^2 - 900]$
 $= 90\ 000 - 100(x - 30)^2$

$a = 90\ 000$
 $b = 100$
 $c = 30$ } _____ 1A

For the method of completing square

All three answers must be correct.

(c) (2 marks)

When $x = 30,$ P is a maximum. _____ 1M+1A

