

**只限教師參閱**

**FOR TEACHERS' USE ONLY**

**香港考試及評核局**

**HONG KONG EXAMINATIONS AND ASSESSMENT AUTHORITY**

**2009年香港中學會考**

**HONG KONG CERTIFICATE OF EDUCATION EXAMINATION 2009**

**數學 試卷一**

**MATHEMATICS PAPER 1**

本評卷參考乃香港考試及評核局專為今年本科考試而編寫，供閱卷員參考之用。閱卷員在完成閱卷工作後，若將本評卷參考提供其任教會考班的本科同事參閱，本局不表反對，但須切記，在任何情況下均不得容許本評卷參考落入學生手中。學生若索閱或求取此等文件，閱卷員/教師應嚴詞拒絕，因學生極可能將評卷參考視為標準答案，以致但知硬背死記，活剝生吞。這種落伍的學習態度，既不符現代教育原則，亦有違考試着重理解能力與運用技巧之旨。因此，本局籲請各閱卷員/教師通力合作，堅守上述原則。

This marking scheme has been prepared by the Hong Kong Examinations and Assessment Authority for markers' reference. The Authority has no objection to markers sharing it, after the completion of marking, with colleagues who are teaching the subject. However, under no circumstances should it be given to students because they are likely to regard it as a set of model answers. Markers/teachers should therefore firmly resist students' requests for access to this document. Our examinations emphasise the testing of understanding, the practical application of knowledge and the use of processing skills. Hence the use of model answers, or anything else which encourages rote memorisation, should be considered outmoded and pedagogically unsound. The Authority is counting on the co-operation of markers/teachers in this regard.

©香港考試及評核局 保留版權  
Hong Kong Examinations and Assessment Authority  
All Rights Reserved 2009



2009-CE-MATH 1-1

**只限教師參閱**

**FOR TEACHERS' USE ONLY**

Solution	Marks	Remarks
1. $\frac{3n-5m}{2} = 4$ $3n-5m = (2)(4)$ $3n = 8+5m$ $n = \frac{8+5m}{3}$	1M 1M 1A	for putting $n$ on one side or equivalent
$\frac{3n-5m}{2} = 4$ $\frac{3n}{2} - \frac{5m}{2} = 4$ $\frac{3n}{2} = 4 + \frac{5m}{2}$ $n = \frac{8}{3} + \frac{5m}{3}$	1M 1M 1A	for putting $n$ on one side or equivalent
	-----(3)	
2. $\frac{x^2}{(x^{-7}y)^3}$ $= \frac{x^2}{x^{-21}y^3}$ $= \frac{x^{2+21}}{y^3}$ $= \frac{x^{23}}{y^3}$	1M 1M 1A -----(3)	for $(ab)^k = a^k b^k$ or $(a^l)^k = a^{lk}$ for $\frac{1}{c^{-m}} = c^m$ or $\frac{c^n}{c^m} = c^{n-m}$
3. (a) $a^2b + ab^2$ $= ab(a+b)$ (b) $a^2b + ab^2 + 7a + 7b$ $= ab(a+b) + 7a + 7b$ $= ab(a+b) + 7(a+b)$ $= (a+b)(ab+7)$	1A 1M 1A -----(3)	or equivalent for using the result of (a) or equivalent

Solution	Marks	Remarks
<p>4. (a) 405.504 = 406 (correct to the nearest integer)</p> <p>(b) 405.504 = 405.50 (correct to 2 decimal places)</p> <p>(c) 405.504 = 410 (correct to 2 significant figures)</p>	<p>1A</p> <p>1A</p> <p>1A</p> <p>----- (3)</p>	
<p>5. The required probability</p> $= \frac{7+21}{7+21+30+53+57+32}$ $= \frac{28}{200}$ $= \frac{7}{50}$	<p>1M + 1M</p> <p>1A</p> <p>----- (3)</p>	<p>1M for numerator + 1M for denominator</p> <p>0.14</p>
<p>6. Let <math>x</math> be the number of stamps owned by John. Then, the number of stamps owned by Mary is <math>(300 - x)</math>.</p> $4x = (300 - x) + 20$ $4x = 320 - x$ $5x = 320$ $x = 64$ <p>Thus, the number of stamps owned by John is 64.</p>	<p>1A</p> <p>1M + 1A</p> <p>1A</p>	<p>pp-1 for any undefined symbol can be absorbed</p>
<p>Let <math>x</math> and <math>y</math> be the numbers of stamps owned by John and Mary respectively.</p> $\begin{cases} x + y = 300 \\ 4x = y + 20 \end{cases}$ <p>So, we have <math>4x = (300 - x) + 20</math>.</p> <p>Solving, we have <math>x = 64</math>.</p> <p>Thus, the number of stamps owned by John is 64.</p>	<p>} 1A + 1A</p> <p>1M</p> <p>1A</p>	<p>pp-1 for any undefined symbol</p> <p>for getting a linear equation in <math>x</math> or <math>y</math> only</p>
	<p>----- (4)</p>	

Solution	Marks	Remarks
7. (a) The required number = $172(1 - 75\%)$ = 43	1M 1A	can be absorbed
(b) The required percentage = $\left(\frac{43}{43+172}\right)(100\%)$ = 20%	1M 1A	accept without 100 %
The required percentage = $\left(1 - \frac{172}{43+172}\right)(100\%)$ = 20%	1M 1A	accept without 100 %
The required percentage = $\left(\frac{1-75\%}{1+(1-75\%)}\right)(100\%)$ = 20%	1M 1A	accept without 100 %
	------(4)	
8. (a) $\angle POQ$ = $213^\circ - 123^\circ$ = $90^\circ$ Thus, $\triangle OPQ$ is a right-angled triangle.	1M 1A	for considering $\angle POQ$ f.t.
(b) $k^2 + 24^2 = 25^2$ $k = 7$ The perimeter of $\triangle OPQ$ = $7 + 24 + 25$ = 56	1M 1A 1M	for using Pythagoras' theorem
	------(5)	
9. (a) The coordinates of $A'$ are $(-1, 4)$ . The coordinates of $B'$ are $(-5, 2)$ .	1A 1A	pp-1 for missing '(' or ')' pp-1 for missing '(' or ')'
(b) The slope of $AB$ = $\frac{2 - (-2)}{5 - (-1)}$ = $\frac{2}{3}$ The slope of $A'B'$ = $\frac{4 - 2}{-1 - (-5)}$ = $\frac{1}{2}$ Since the slope of $AB$ and the slope of $A'B'$ are different, $AB$ is not parallel to $A'B'$ .	1M 1A 1A	either one either one f.t.
	------(5)	

Solution	Marks	Remarks
10. (a) The median = 26 words per minute	1A	u-1 for missing unit
The range = 39 - 12 = 27 words per minute	1A	u-1 for missing unit
The inter-quartile range = 35 - 21 = 14 words per minute	1A	u-1 for missing unit
----- (3)		
(b) (i) The inter-quartile range after the training = 47 - 35 = 12 words per minute	1M	for considering inter-quartile range
The inter-quartile range of the distribution of typing speed after the training is less than that before the training. Thus, the distribution after the training is not more dispersed than that before the training.	1A	f.t.
The range after the training = 52 - 27 = 25 words per minute	1M	for considering range
The range of the distribution of typing speed after the training is less than that before the training. Thus, the distribution after the training is not more dispersed than that before the training.	1A	f.t.
(ii) Note that the median of the distribution of the typing speed after the training is 40 words per minute which is greater than the maximum of the distribution of the typing speed before the training. Thus, the claim is agreed.	1M 1A	f.t.
Note that the minimum of the distribution of the typing speed after the training is 27 words per minute which is greater than the median of the distribution of the typing speed before the training. Thus, the claim is agreed.	1M 1A	f.t.
----- (4)		

Solution	Marks	Remarks
11. <b>Marking Schemes for (a) and (b)(i):</b>		
<b>Case 1</b> Any correct proof with correct reasons.	3	
<b>Case 2</b> Any correct proof without reasons.	2	
<b>Case 3</b> Incomplete proof with any one correct step and one correct reason.	1	
<p>(a) In <math>\triangle ABC</math> and <math>\triangle AED</math>,</p> <p><math>AC = AD</math> ( given )</p> <p><math>BC = ED</math> ( given )</p> <p><math>\angle ACB + \angle BCE = \angle ADE + \angle CAD</math> ( ext. <math>\angle</math> of <math>\triangle</math> )</p> <p><math>\angle BCE = \angle CAD</math> ( given )</p> <p><math>\angle ACB = \angle ADE</math></p> <p><math>\triangle ABC \cong \triangle AED</math> ( SAS )</p>		<p>[已知]</p> <p>[已知]</p> <p>[<math>\triangle</math>的外角]</p> <p>[已知]</p>
	----- (3)	
<p>(b) (i) In <math>\triangle ABF</math> and <math>\triangle DEA</math>,</p> <p><math>\angle AFB = \angle DAE</math> (alt. <math>\angle</math>s, <math>AD \parallel BC</math>)</p> <p><math>\angle FBA = \angle AED</math> ( by (a) )</p> <p><math>\angle FAB = \angle ADE</math> (<math>\angle</math> sum of <math>\triangle</math> )</p> <p><math>\triangle ABF \sim \triangle DEA</math> ( AAA )</p>		<p>[[內]錯角, <math>AD \parallel BC</math>]</p> <p>[由(a)]</p> <p>[<math>\triangle</math>內角和]</p> <p>(AA) (equiangular)</p> <p>[等角]</p>
<p>(ii) <math>\triangle CBA</math> and <math>\triangle CEF</math> are similar to <math>\triangle ABF</math>.</p>	1A + 1A	
	----- (5)	

Solution	Marks	Remarks
12. (a) (i) The equation of the axis of symmetry is $x = 11$ .	1A	or equivalent
(ii) The coordinates of $R$ are $(11, 23)$ .	1A	pp-1 for missing '(' or ')'
-----(2)		
(b) (i) $-2(x-11)^2 + 23 = 5$ $(x-11)^2 = 9$ $x-11 = 3$ or $x-11 = -3$ $x = 14$ or $x = 8$	1M  1A	
$-2(x-11)^2 + 23 = 5$ $2x^2 - 44x + 224 = 0$ $x^2 - 22x + 112 = 0$ $(x-8)(x-14) = 0$ $x = 8 \text{ or } x = 14$	1M  1A	
<p>The distance between <math>P</math> and <math>Q</math></p> $= 14 - 8$ $= 6$	1M 1A	can be absorbed
(ii) The area of $\triangle PQR$ $= \frac{(6)(23-5)}{2}$ $= 54$	1M 1A	<div style="text-align: center;">either one</div>
<p>The area of <math>\triangle PSQ</math></p> $= \frac{(6)(5)}{2}$ $= 15$		<div style="text-align: center;">either one</div>
<p>The area of the quadrilateral <math>PRQS</math></p> $= 54 + 15$ $= 69$	1A	
<p>The area of the quadrilateral <math>PRQS</math></p> $= \frac{(6)(23)}{2}$ $= 69$	1M + 1A 1A	
-----(7)		

Solution	Marks	Remarks
<p>13. (a) (i) The capacity of the container</p> $= \frac{1}{3}\pi(12)^2(18)$ $= 864\pi \text{ cm}^3$ <p>(ii) Let <math>r</math> cm be the base radius of the lower part of the container.</p> <p>Then, we have <math>\frac{18-6}{18} = \frac{r}{12}</math>.</p> <p>Solving, we have <math>r = 8</math>.</p> <p>The volume of frustum</p> $= 864\pi - \frac{1}{3}\pi(8)^2(18-6)$ $= 864\pi - 256\pi$ $= 608\pi \text{ cm}^3$	<p>1M</p> <p>1A</p> <p>1M</p> <p>1A</p>	<p>u-1 for missing unit</p> <p>u-1 for missing unit</p>
<p>Let <math>V \text{ cm}^3</math> be the volume of the frustum.</p> <p>Then, we have <math>\frac{864\pi - V}{864\pi} = \left(\frac{18-6}{18}\right)^3</math>.</p> <p>So, we have <math>\frac{864\pi - V}{864\pi} = \frac{8}{27}</math>.</p> <p>Solving, we have <math>V = 608\pi</math>.</p> <p>Thus, the volume of the frustum is <math>608\pi \text{ cm}^3</math>.</p>	<p>1M</p> <p>1A</p>	<p>u-1 for missing unit</p>
------(4)		
<p>(b) (i) The capacity of the lower part of the vessel</p> $= \pi(8)^2(10)$ $= 640\pi \text{ cm}^3$ <p>Let <math>h</math> cm be the depth of water in the vessel.</p> $\left(\frac{h-10+(18-6)}{18}\right)^3 = \frac{(884\pi - 640\pi) + (864\pi - 608\pi)}{864\pi}$ $\left(\frac{h+2}{18}\right)^3 = \frac{125}{216}$ $\frac{h+2}{18} = \frac{5}{6}$ $h = 13$ <p>Thus, the depth of water in the vessel is 13 cm.</p>	<p>1M</p> <p>1M</p> <p>1A</p>	<p>u-1 for missing unit</p>
<p>(ii) The volume not occupied by water in the vessel</p> $= 608\pi + 640\pi - 884\pi$ $= 364\pi \text{ cm}^3$ <p>Note that the volume of metal is <math>1000 \text{ cm}^3</math>.</p> <p>So, the volume of metal is less than <math>364\pi \text{ cm}^3</math>.</p> <p>Thus, the water will not overflow.</p>	<p>1M</p> <p>1A</p>	<p>for comparison f.t.</p>
------(5)		





Solution	Marks	Remarks
<p>15. (a) (i) The taxi fare  <math>= 30 + \frac{x-2}{0.2}(2.4)</math>  <math>= 30 + 12(x-2)</math>  <math>= \\$ (6 + 12x)</math></p> <p>(ii) If <math>x</math> is not a multiple of 0.2, then the distance used for calculating the taxi fare is the estimated distance after rounding up to the nearest 0.2 km instead of the actual distance travelled.                      Thus, the taxi fare is not <math>\\$ (6 + 12x)</math>.</p>	<p>1</p> <p>1A</p> <p>----- (2)</p>	<p>f.t.</p>
<p>(b) Note that the distance used for calculating the taxi fare is 3.2 km .                      The taxi fare  <math>= 6 + (12)(3.2)</math>  <math>= \\$ 44.4</math></p>	<p>1A</p> <p>1M</p> <p>1A</p> <p>----- (3)</p>	<p>can be absorbed</p>
<p>(c) The taxi fare for the 2nd journey  <math>= 6 + (12)(3.6)</math>  <math>= \\$ 49.2</math></p> <p>The distance covered by the 99th journey  <math>= 3.1 + (99-1)(0.5)</math>  <math>= 52.1</math> km</p> <p>The taxi fare for the 99th journey  <math>= 6 + (12)(52.2)</math>  <math>= \\$ 632.4</math></p> <p>The total taxi fare  <math>= 44.4 + 49.2 + 56.4 + 61.2 + \dots + 625.2 + 632.4</math>  <math>= (44.4 + 56.4 + \dots + 632.4) + (49.2 + 61.2 + \dots + 625.2)</math>  <math>= \frac{(44.4 + 632.4)(50)}{2} + \frac{(49.2 + 625.2)(49)}{2}</math>  <math>= 16\ 920 + 16\ 522.8</math>  <math>= \\$ 33\ 442.8</math>  <math>&gt; \\$ 33\ 000</math>                      Thus, the claim is incorrect.</p>	<p>1A</p> <p>1M</p> <p>1M+1A+1A</p> <p>1A</p> <p>----- (6)</p>	<p>can be absorbed</p> <p>for splitting into two arithmetic sequences</p> <p>1M for summing either arithmetic sequence + 1A for correct number of terms in each series</p> <p>f.t.</p>
<p>The taxi fare for the 2nd journey  <math>= 6 + (12)(3.6)</math>  <math>= \\$ 49.2</math></p> <p>The total taxi fare  <math>= 44.4 + 49.2 + 56.4 + 61.2 + \dots + 625.2 + 632.4</math>  <math>= (44.4 + 56.4 + \dots + 632.4) + (49.2 + 61.2 + \dots + 625.2)</math>  <math>= \frac{((2)(44.4) + (50-1)(12))(50)}{2} + \frac{((2)(49.2) + (49-1)(12))(49)}{2}</math>  <math>= 16\ 920 + 16\ 522.8</math>  <math>= \\$ 33\ 442.8</math>  <math>&gt; \\$ 33\ 000</math>                      Thus, the claim is incorrect.</p>	<p>1A</p> <p>1M</p> <p>1M+1A+1A</p> <p>1A</p> <p>----- (6)</p>	<p>can be absorbed</p> <p>for splitting into two arithmetic sequences</p> <p>1M for summing either arithmetic sequence + 1A for correct number of terms in each series</p> <p>f.t.</p>

Solution	Marks	Remarks
<p>16 (a) (i) The slope of <math>L_1</math></p> $= \frac{24-16}{12-8}$ $= 2$ <p>The equation of <math>L_1</math> is</p> $y-16 = 2(x-8)$ $2x-y=0$ <p>The slope of <math>L_2</math></p> $= \frac{-1}{2}$ <p>The equation of <math>L_2</math> is</p> $y-24 = \frac{-1}{2}(x-12)$ $x+2y-60=0$ <p>(ii) The system of inequalities is</p> $\begin{cases} x \geq 8 \\ y \geq 10 \\ 2x \geq y \\ x+2y \leq 60 \end{cases}$	<p>1M 1A</p> <p>1A</p> <p>1A+1A+1A</p> <p>(6)</p>	<p>or equivalent</p> <p>either one</p> <p>or equivalent</p> <p>or equivalent</p>
<p>(b) Let <math>x</math> and <math>y</math> be the numbers of square tables and round tables placed respectively.</p> <p>Now, the constraints are <math>x \geq 8</math>, <math>y \geq 10</math>, <math>2x \geq y</math> and <math>x+2y \leq 60</math>.</p> <p>Denote the total profit on the dining tables by <math>\\$P</math>.</p> <p>Then, we have <math>P = 4\,000x + 6\,000y</math>.</p> <p>Note that the vertices of the shaded region are <math>(12, 24)</math>, <math>(40, 10)</math>, <math>(8, 10)</math> and <math>(8, 16)</math>.</p> <p>At <math>(12, 24)</math>, we have <math>P = (4\,000)(12) + (6\,000)(24) = 192\,000</math>.</p> <p>At <math>(40, 10)</math>, we have <math>P = (4\,000)(40) + (6\,000)(10) = 220\,000</math>.</p> <p>At <math>(8, 10)</math>, we have <math>P = (4\,000)(8) + (6\,000)(10) = 92\,000</math>.</p> <p>At <math>(8, 16)</math>, we have <math>P = (4\,000)(8) + (6\,000)(16) = 128\,000</math>.</p> <p>So, the greatest value of <math>P</math> is <math>220\,000</math>.</p> <p>Thus, the claim is disagreed.</p>	<p>1A</p> <p>1M + 1M</p> <p>1A 1A</p>	<p>1M for testing a point + 1M for testing four points</p> <p>f.t.</p>
<p>Let <math>x</math> and <math>y</math> be the numbers of square tables and round tables placed respectively.</p> <p>Now, the constraints are <math>x \geq 8</math>, <math>y \geq 10</math>, <math>2x \geq y</math> and <math>x+2y \leq 60</math>.</p> <p>Denote the total profit on the dining tables by <math>\\$P</math>.</p> <p>Then, we have <math>P = 4\,000x + 6\,000y</math>.</p> <p>Draw the straight line <math>2x+3y=k</math> on Figure 5, where <math>k</math> is a constant.</p> <p>It is found that <math>P</math> attains its greatest value at <math>(40, 10)</math>.</p> <p>So, the greatest value of <math>P</math> is <math>220\,000</math>.</p> <p>Thus, the claim is disagreed.</p>	<p>1A</p> <p>1M + 1M</p> <p>1A 1A</p>	<p>1M for sliding straight line + 1M for straight line with negative slope</p> <p>f.t.</p>
	<p>(5)</p>	



Solution	Marks	Remarks
<p>(b) <math>DE</math>  <math>= \sqrt{CD^2 - CE^2}</math>  <math>\approx \sqrt{22.30713539^2 - 7^2}</math>  <math>\approx 21.1803751</math> cm</p> <p>Let <math>d</math> be the perpendicular distance from <math>E</math> to <math>CD</math>.                      Then, we have <math>\frac{d(CD)}{2} = \frac{(CE)(DE)}{2}</math>.                      Therefore, we have <math>d = \frac{(CE)(DE)}{CD}</math>.</p> <p>The perpendicular distance from <math>E</math> to <math>CD</math>  <math>= \frac{(CE)(DE)}{CD}</math>  <math>\approx \frac{(7)(21.1803751)}{22.30713539}</math>  <math>\approx 6.646421565</math> cm</p> <p>So, the perpendicular distance from <math>E</math> to <math>CD</math> and the shortest distance from <math>E</math> to the horizontal ground are different.                      Thus, the claim is disagreed.</p>	<p>1M</p> <p>1A</p>	<p>accept equating sine ratios</p> <p>f.t.</p>
<p><math>DE</math>  <math>= \sqrt{CD^2 - CE^2}</math>  <math>\approx \sqrt{22.30713539^2 - 7^2}</math>  <math>\approx 21.1803751</math> cm</p> <p>Let <math>\theta</math> be the angle between <math>DE</math> and the horizontal ground.  <math>\sin \theta = \frac{\text{The shortest distance from } E \text{ to the horizontal ground}}{DE}</math>  <math>\sin \theta \approx \frac{6.598354429}{21.1803751}</math>  <math>\theta \approx 18.1515512^\circ</math></p> <p><math>\sin \angle CDE = \frac{CE}{CD}</math>  <math>\sin \angle CDE \approx \frac{7}{22.30713539}</math>  <math>\angle CDE \approx 18.28844266^\circ</math></p> <p>So, <math>\angle CDE</math> and the angle between <math>DE</math> and the horizontal ground are different.                      Thus, the claim is disagreed.</p>	<p>1M</p> <p>1A</p>	<p>for identifying the required angle</p> <p>f.t.</p>
	<p>------(2)</p>	