

1980 A-Level Pure Mathematics Paper I

1. Let V be the set of all 3×1 real matrices. For any $\underline{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$, $\underline{y} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$ in V and any real number λ ,

we define $\underline{x} + \underline{y} = \begin{pmatrix} x_1 + y_1 \\ x_2 + y_2 \\ x_3 + y_3 \end{pmatrix}$, $\lambda \underline{x} = \begin{pmatrix} \lambda x_1 \\ \lambda x_2 \\ \lambda x_3 \end{pmatrix}$. It is known that under this addition and scalar multiplication,

V forms a real vector space with zero vector $\underline{0} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$.

(a) For a given 3×3 real matrix A , let $E = \{\underline{x} \in V : A\underline{x} = \underline{0}\}$.

(i) Show that E forms a vector subspace of V .

(ii) For \underline{b} in V , suppose we have \underline{p} in V such that $A\underline{p} = \underline{b}$. Show that, for any \underline{y} in V ,

$A\underline{y} = \underline{b}$ if and only if $\underline{y} = \underline{p} + \underline{x}$ for some \underline{x} in E .

(b) (i) Find all solutions to
$$\begin{cases} x - y - z = 0 \\ 10x + 5y - 4z = 0 \\ 5x + 5y - z = 0 \end{cases}$$

(ii) Suppose $x = \frac{1}{2}$, $y = \frac{4}{3}$, $z = \sqrt{2}$ is a solution to the system of equations

$$\begin{cases} x - y - z = b_1 \\ 10x + 5y - 4z = b_2 \\ 5x + 5y - z = b_3 \end{cases}.$$

Find all solutions to the system.

(1980)

2. Let F denote the set of all positive-valued continuous functions on the set \mathbb{R} of all real numbers. For any $f, g \in F$, define $f * g$ by $(f * g)(x) = f(x)g(x) \quad \forall x \in \mathbb{R}$. It is known that F forms a group under the operation $*$. The identity I of this group and the inverse g of $f \in F$ are given respectively by

$$I(x) = 1 \quad \forall x \in \mathbb{R}, \quad g(x) = \frac{1}{f(x)} \quad \forall x \in \mathbb{R}.$$

Define a relation \sim in F as follows:

For $f, g \in F$, $f \sim g$ if there are polynomials p, q in F such that $p * f = q * g$.

(a) Show that \sim is an equivalence relation on F .

(b) Let f / \sim be the equivalence class of f with respect to \sim , and let F / \sim be the quotient set consisting of all these equivalence classes. For any $f / \sim, g / \sim \in F / \sim$, define $f / \sim \otimes g / \sim$ to be $(f * g) / \sim$.

(i) Show that \otimes is well defined on F / \sim , i.e., if $f / \sim = f_1 / \sim$ and

$$g / \sim = g_1 / \sim, \text{ then } f / \sim \otimes g / \sim = f_1 / \sim \otimes g_1 / \sim.$$

(ii) Show that F / \sim forms a group under \otimes .

(1980)

3. (a) If $x > 0$ and p is a positive integer, show that $\frac{x^{p+1} - 1}{p+1} \geq \frac{x^p - 1}{p}$, and that the equality holds only if $x = 1$.

(b) Let x_1, x_2, \dots, x_n be positive numbers and $\sum_{i=1}^n x_i \geq n$.

(i) Show that, for any positive integer m , $\sum_{i=1}^n x_i^m \geq n$.

(ii) If $\sum_{i=1}^n x_i^m = n$ for some integer m greater than one, show that $x_1 = x_2 = \dots = x_n = 1$.

(c) Using (b), or otherwise, show that, for any positive numbers y_1, y_2, \dots, y_n , and positive integer m ,

$$\frac{y_1^m + y_2^m + \dots + y_n^m}{n} \geq \left(\frac{y_1 + y_2 + \dots + y_n}{n} \right)^m \quad \text{and that the equality holds only when } m=1 \text{ or } y_1 = y_2 = \dots = y_n. \quad (1980)$$

4. (a) The terms of a sequence y_1, y_2, y_3, \dots satisfy the relation $y_k = Ay_{k-1} + B$ ($k \geq 2$) where A, B are constants independent of k and $A \neq 1$. Guess an expression for y_k ($k \geq 2$) in terms of y_1, A, B and k and prove it.

(b) The terms of a sequence x_0, x_1, x_2, \dots satisfy the relation $x_k = (a+b)x_{k-1} - abx_{k-2}$ ($k \geq 2$), where a, b are non-zero constants independent of k and $a \neq b$.

(i) Express $x_k - ax_{k-1}$ ($k \geq 2$) in terms of $(x_1 - ax_0), b$ and k .

(ii) Using (a) or otherwise, express x_k ($k \geq 2$) in terms of x_0, x_1, a, b and k .

(c) If the terms of the sequence x_0, x_1, x_2, \dots satisfy the relation $x_k = \frac{1}{3}x_{k-1} + \frac{2}{3}x_{k-2}$ ($k \geq 2$), express $\lim_{k \rightarrow \infty} x_k$ in terms of x_0 and x_1 . (1980)

5. (a) (i) Let $\omega^3 = 1$ and $\omega \neq 1$. Show that the expression $x^3 - 3uvx - (u^3 + v^3) = 0$ can be factorized as $(x - u - v)(x - \omega u - \omega^2 v)(x - \omega^2 u - \omega v)$

(ii) Find a solution to the following system of equations

$$\begin{cases} u^3 + v^3 = 6 \\ uv = 2 \end{cases}$$

Hence, or otherwise, find the roots of the equations $x^3 - 6x - 6 = 0$

(b) Given an equation $x^3 + px + q = 0$ (*)

(i) Show that, if (*) has a multiple root, then $27q^2 + 4p^3 = 0$

(ii) Using the method indicated in (a) (ii), or otherwise, show that, if $27q^2 + 4p^3 = 0$, then (*) has a multiple root. (1980)

6. Let a, b be real numbers such that $a < b$ and let m, n be positive integers.

(a) If for all real numbers x, u , $[(1+u)x - (au+b)]^{m+n} = \sum_{k=0}^{m+n} A_k(x)u^k$ (*)

show that $A_k(x) = C_k^{m+n} (x-a)^k (x-b)^{m+n-k}$ for $k = 0, 1, \dots, m+n$,

where C_k^{m+n} is the coefficient of t^k in the expansion of $(1+t)^{m+n}$.

(b) By integrating both sides of (*) with respect to x , or otherwise, calculate $\int_a^b (x-a)^m (x-b)^n dx$.

(c) By differentiating both sides of (*) with respect to x , or otherwise, find $\frac{d^r}{dx^r} \{(x-a)^m (x-b)^n\}$ at $x=a$, where r is a positive integer. (1980)

7. Let C be the set of complex numbers. A function $f: C \rightarrow C$ is said to be an isometry if it preserves distance, that is, if $|f(z_1) - f(z_2)| = |z_1 - z_2|$ for all $z_1, z_2 \in C$.

(a) If f is an isometry, show that $g(z) = \frac{f(z) - f(0)}{f(1) - f(0)}$ is an isometry satisfying $g(1) = 1$ and $g(0) = 0$.

(b) If g is an isometry satisfying $g(1) = 1$, $g(0) = 0$, show that

(i) the real parts of $g(z)$ and z are equal for all $z \in C$,

(ii) $g(i) = i$ or $-i$.

(c) If g is an isometry satisfying $g(1) = 1$, $g(0) = 0$ and $g(i) = i$ (respectively $-i$), show that $g(z) = z$ (respectively \bar{z}) for all $z \in C$.

(d) Show that any isometry f has the form $f(z) = az + b$ or $f(z) = a\bar{z} + b$ with a and b constant and $|a| = 1$. (1980)

8. N balls are distributed randomly among n cells. Each of the n^N possible distributions has probability n^{-N} .

(a) (i) Calculate the probability P_k that a given cell contains exactly k balls.

(ii) Show that the most probable number k_0 satisfies the inequality $\frac{N-n+1}{n} \leq k_0 \leq \frac{N+1}{n}$.

(iii) Compute the mean number $\sum_{k=0}^N kP_k$ of balls in a given cell and show that it can differ from k_0 by at most one.

(b) Let $A(N, n)$ be the number of distributions leaving none of the cells empty. Show that

$A(N, n+1) = \sum_{k=1}^N C_k^N A(N-k, n)$, where C_k^N is the coefficient of t^k in the expansion of

$(1+t)^N$. Hence show by mathematical induction (on n), or otherwise, that

$$A(N, n) = \sum_{j=0}^n (-1)^j C_j^n (n-j)^N. \quad (1980)$$

1980 A-Level Pure Mathematics Paper II

1. Let $P(t) = (x(t), y(t))$ be a point on the unit circle with parametric equations
 $x(t) = \frac{1-t^2}{1+t^2}$, $y(t) = \frac{2t}{1+t^2}$, Q be the point $(a, 0)$, $0 < a < 1$. An arbitrary line of slope m passing through Q cuts the circle at the points $R = P(t_1)$ and $S = P(t_2)$. Let T be the point where RO meets the line through Q parallel to SO , where O is the origin.
- (a) Show that $t_1 t_2 = \frac{a-1}{a+1}$, $t_1 + t_2 = \frac{-2}{m(a+1)}$.
- (b) Express the coordinates of T in terms of a , t_1 , t_2
- (c) Verify that the locus of T is an ellipse with equation $(1-a^2)(x - \frac{a}{2})^2 + y^2 = C$, where C is a constant. What is C ?
 (1980)

2. In a 3-dimensional space with a Cartesian coordinate system, two lines l_1 and l_2 are given by the pairs of equations:

$$l_1: \begin{cases} x + 2y + 3z - 3 = 0 \\ x + 2y + 2z - 4 = 0 \end{cases}, \quad l_2: \begin{cases} x + y + z - 1 = 0 \\ 2x + 3y + 5z - 2 = 0 \end{cases}.$$

- (a) Let P_α be the plane $\alpha(x + 2y + 3z - 3) + (x + 2y + 2z - 4) = 0$ and Q_β be the plane $\beta(x + y + z - 1) + (2x + 3y + 5z - 2) = 0$. Show that P_α is parallel to Q_β if and only if there exists $m \neq 0$ such that

$$\begin{pmatrix} 1 & -m & 1-2m \\ 2 & -m & 2-3m \\ 3 & -m & 2-5m \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \dots\dots\dots (*)$$


- (b) Find the value of m for which there are numbers α and β satisfying (*) in (a). Hence find the equations of the two parallel planes M_1 and M_2 containing l_1 and l_2 respectively.
- (c) Find the equation of the plane N containing l_1 and perpendicular to M_2 .
- (d) Let l_1' be the projection of l_1 on M_2 (i.e. l_1' is the intersection of N and M_2). Find its equation.
 (1980)

3. (a) Let $f(z) = \sum_{k=0}^n a_k z^k$ be an n^{th} degree polynomial in the complex variable z with real coefficients. Show that
- (i) $|f(z)|^2 = \sum_{k=0}^n \sum_{j=0}^n r^{k+j} a_k a_j \cos(k-j)\theta$, where $z = r(\cos\theta + i \sin\theta)$,
- (ii) $\frac{1}{2\pi} \int_0^{2\pi} |f(\cos\theta + i \sin\theta)|^2 d\theta = \sum_{k=0}^n a_k^2$.
- (b) If C_k^n is the coefficient of t^k in the binomial expansion of $(1+t)^n$, show that


$$\sum_{k=0}^n \binom{n}{k}^2 = \frac{2^n}{\pi} \int_0^\pi (1 + \cos\theta)^n d\theta = \frac{2^{2n+1}}{\pi} \int_0^{\frac{\pi}{2}} (\cos t)^{2n} dt.$$

(1980)

4. (a) A right circular cone is inscribed in a sphere of radius a as shown in the figure. Determine the height of the cone if it is to have maximum volume.



- (b) Two points A and B lie on the circumference of a circle with centre O and radius a such that $\angle AOB = \frac{2}{3}\pi$. X is a point on AO produced with $OX = x$; P is a point on arc AB with $\angle AOP = \theta$. For each $x > 0$, let $g(x) = \int_0^{\frac{2\pi}{3}} \frac{a \sin\theta}{r(x, \theta)} d\theta$, where $r(x, \theta)$ is the distance between P and X .



(i) Show that
$$g(x) = \frac{3a}{x + a + (a^2 + x^2 - ax)^{\frac{1}{2}}}.$$

(ii) Prove that on $(0, \infty)$, $0 < g(x) < \frac{3}{2}$.

(1980)

5. (a) Let f and g be two continuous functions defined on the real line \mathbb{R} and let $x_0 \in \mathbb{R}$, show that if $f(x) = g(x)$ for all $x \in \mathbb{R} \setminus \{x_0\}$, then $f(x_0) = g(x_0)$.

- (b) If a real polynomial $p(x)$ can be written as $p(x) = (x - x_0)^m q(x)$ for some positive integer m and polynomial $q(x)$ with $q(x_0) \neq 0$, show that the expression is unique, that is, if k is a positive integer and $h(x)$ is a polynomial with $h(x_0) \neq 0$ such that $p(x) = (x - x_0)^k h(x)$, then $m = k$ and $q(x) = h(x)$ for all $x \in \mathbb{R}$.

- (c) Let $p(x)$ be a real polynomial. Show that for any positive integer k , x_0 is a root of multiplicity $k + 1$ of the equation $p(x) = 0$ if and only if $p(x_0) = 0$ and x_0 is a root of multiplicity k of $p'(x) = 0$, where $p'(x)$ denotes the derivative of $p(x)$.
(1980)

6. (a) Let f and g be real-valued functions defined on the real line \mathbf{R} and possess the following properties :

(1) $f(x + y) = f(x)g(y) + f(y)g(x)$ for all $x, y \in \mathbf{R}$,

(2) $f(0) = 0, f'(0) = 1, g(0) = 1, g'(0) = 0$

Show that $f'(x) = g(x)$ for all $x \in \mathbf{R}$.

(b) Let $f(x)$ be a function with continuous first and second derivatives on $[0, 1]$ and $f(0) = f(1) = 0$.

(i) Show that $\int_0^1 f(x)f''(x)dx \leq 0$, where the equality sign holds only if $f(x) = 0$ for all x in $[0, 1]$.

(ii) Suppose, in addition, $\int_0^1 [f(x)]^2 dx = 1$. Show that $\int_0^1 xf(x)f'(x)dx = -\frac{1}{2}$

(1980)

7. (a) It is known that, for any integer k , $\int_{-\pi}^{\pi} \sin kx dx = 0$ and $\int_{-\pi}^{\pi} \cos kx dx = \begin{cases} 2\pi & \text{if } k=0 \\ 0 & \text{if } k \neq 0 \end{cases}$

Using the above results, show that if m, n are positive integers,

(i) $\int_{-\pi}^{\pi} \sin mx \cos nx dx = 0,$ (ii)

$$\int_{-\pi}^{\pi} \sin mx \sin nx dx = \begin{cases} \pi & \text{if } m=n, \\ 0 & \text{if } m \neq n, \end{cases}$$

(iii) $\int_{-\pi}^{\pi} \cos mx \cos nx dx = \begin{cases} \pi & \text{if } m=n, \\ 0 & \text{if } m \neq n. \end{cases}$

(b) Define $\phi_i(x) = \begin{cases} \frac{1}{\sqrt{2\pi}} & \text{if } i=0, \\ \frac{\cos mx}{\sqrt{\pi}} & \text{if } i=2m-1, \\ \frac{\sin mx}{\sqrt{\pi}} & \text{if } i=2m, \end{cases}$,

where $m = 1, 2, 3, \dots$

Let f be a continuous real-valued function defined on $[-\pi, \pi]$ and let α_i be real constants.

(i) Prove that, for each integer $N \geq 0$,

$$\int_{-\pi}^{\pi} [f(x) - \sum_{i=0}^N \alpha_i \phi_i(x)]^2 dx = \int_{-\pi}^{\pi} [f(x)]^2 dx + \sum_{i=0}^N \alpha_i^2 - 2 \sum_{i=0}^N \alpha_i p_i,$$

where $p_i = \int_{-\pi}^{\pi} f(x)\phi_i(x)dx$. Hence prove that $\int_{-\pi}^{\pi} [f(x) - \sum_{i=0}^N \alpha_i \phi_i(x)]^2 dx$

attains its least value for varying α_i when $\alpha_i = p_i$ for each i .

(ii) Show that, for any integer $M \geq 1$, $\sum_{i=0}^{2M} p_i^2 \leq \int_{-\pi}^{\pi} [f(x)]^2 dx$.

(1980)

8. Let Γ and Γ' be two Cartesian coordinate systems on a plane and with the same origin, where Γ' is obtained from Γ by a rotation through an angle θ . If (x, y) and (x', y') are the coordinates of an arbitrary point P with respect to Γ and Γ' respectively, then it is known that $x' = kx + hy$, $y' = -hx + ky$, where $k = \cos\theta$ and $h = \sin\theta$.

(a) The general equation of a conic section in the coordinate system Γ is given by

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0.$$

(i) Show that the same conic section is represented in the coordinate system Γ' by

$$A'x'^2 + B'x'y' + C'y'^2 + D'x' + E'y' + F' = 0,$$

$$\text{where } A' = Ak^2 + Bkh + Ch^2,$$

$$B' = 2kh(C - A) + B(k^2 - h^2),$$

$$C' = Ah^2 - Bhk + Ck^2,$$

$$D' = Dk + Eh,$$

$$E' = Ek - Dh, \quad F' = F.$$

(ii) Show that $4A'C' - B'^2 = 4AC - B^2$.

(iii) Show that, by choosing a suitable angle θ of rotation, the coefficient B' can be made to vanish.

(b) By a suitable rotation followed by a translation if necessary, bring

the equation of the conic section $x^2 - 2xy + y^2 + \frac{7}{2}x - \frac{1}{2}y + \frac{11}{2} = 0$ into the standard form.

Write down the equation of its line of symmetry in the original coordinate system.

(1980)