

(a) $A \cdot I \rightarrow \det(A^3) = \det I = 1 \Rightarrow |A|^3 = |A \cdot A \cdot A| = |A||A||A|$
 $\rightarrow (\det A)^3 = 1 \Rightarrow \det A = 1$ (as it is real) $= |A|^3$

(b) (i) $B^2 + B + I = 0 \Rightarrow B^2 = -(B+I)$ or $\Rightarrow (B^2+B) = -I$
 $B(B^2+B+I) = 0 \Rightarrow B^3 = -I$
 $B^3 = I$
 $B^2 \cdot B = B \cdot B^2 = I$

(ii) $I + B + B^2 + \dots + B^{100}$
 $= (I+B+B^2) + B^3(1+B+B^2) + \dots + B^{96}(1+B+B^2) + B^{99} + B^{100}$
 $= I + B$ (or $-B^2$) as $B^3 = -I$

Let $B = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$

$B^2 = -B - I$
 $\Rightarrow \frac{1}{\det B} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} = \begin{pmatrix} -(a+1) & -b \\ -c & -(d+1) \end{pmatrix}$

By (a), $\det B = 1$ as $B^3 = I$.
 $\begin{pmatrix} d & -b \\ -c & a \end{pmatrix} = \begin{pmatrix} -(a+1) & -b \\ -c & -(d+1) \end{pmatrix}$
 $d = -(a+1)$ and $a = -(d+1)$
 i.e. $a + d = -1$

(c) Let $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ be a matrix with integral entries such that $M^2 + IM + I = 0$.

By (b), $M \neq I$, $M^2 \neq I$ and $a + d = -1$.

Further $ad - bc = 1$ by (a).
 Putting $a = 0$, then $d = -1$
 Putting $b = 1$, then $c = -1$

Note that $M = \begin{pmatrix} a & b \\ \frac{-1-a-a^2}{b} & -1-a \end{pmatrix}$

On checking, $M = \begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}$ satisfies $M^2 + IM + I = 0$.

3 possible matrices satisfy $M^2 + IM + I = 0$

2(a) (i) $|z|^2 = z\bar{z} = z \frac{a+ib}{1-c}$
 $= \frac{a+ib}{1-c} \cdot \frac{a-ib}{1-c}$
 $= \frac{a^2+b^2}{(1-c)^2}$
 $= \frac{1-c^2}{(1-c)^2}$
 $= \frac{1+c}{1-c}$

(ii) $\frac{1+c}{1-c} = z\bar{z} \Rightarrow 1+c = z\bar{z}(1-c)$
 $\Rightarrow c(1+z\bar{z}) = z\bar{z} - 1$
 $\Rightarrow c = \frac{z\bar{z} - 1}{z\bar{z} + 1}$ ($z\bar{z} + 1 \neq 0$)

(iii) $z + \bar{z} = \frac{a+ib}{1-c} + \frac{a-ib}{1-c}$
 $= \frac{2a}{1-c}$
 $= \frac{2a}{1 - \frac{z\bar{z} - 1}{z\bar{z} + 1}}$
 $= a(z\bar{z} + 1)$

$a = \frac{z + \bar{z}}{z\bar{z} + 1}$
 $z - \bar{z} = \frac{2bi}{1-c}$
 $= \frac{2bi}{1 - \frac{z\bar{z} - 1}{z\bar{z} + 1}}$
 $= 2ib(z\bar{z} + 1)$

$b = \frac{i(z\bar{z} - 1)}{z\bar{z} + 1}$ (or $\frac{a - \bar{z}}{i(z\bar{z} + 1)}$)

$a = \frac{z + \bar{z}}{z\bar{z} + 1}$
 $z - \bar{z} = \frac{2bi}{1-c}$
 $= \frac{2bi}{1 - \frac{z\bar{z} - 1}{z\bar{z} + 1}}$
 $= 2ib(z\bar{z} + 1)$

$b = \frac{i(z\bar{z} - 1)}{z\bar{z} + 1}$ (or $\frac{a - \bar{z}}{i(z\bar{z} + 1)}$)

$a = \frac{z + \bar{z}}{z\bar{z} + 1}$
 $z - \bar{z} = \frac{2bi}{1-c}$
 $= \frac{2bi}{1 - \frac{z\bar{z} - 1}{z\bar{z} + 1}}$
 $= 2ib(z\bar{z} + 1)$

$b = \frac{i(z\bar{z} - 1)}{z\bar{z} + 1}$ (or $\frac{a - \bar{z}}{i(z\bar{z} + 1)}$)



2(a) $\frac{dx}{dt} = 2at, \frac{dy}{dt} = 2a$
 Slope of tangent at $(at^2, 2at)$ ($t \neq 0$) is $\frac{dy}{dx} = \frac{2a}{2at} = \frac{1}{t}$

II

1

$\frac{1}{2}$

1

1

1

$\frac{1}{4}$

1

1

1

1

1

1

1

$\frac{1}{8}$

Equation of the normal is $y - 2at = -t(x - at^2)$
 i.e. $tx + y - (at^3 + 2at) = 0$ (which also holds for $t = 0$)

(b) The equation of the normal at t_1 is

$t_1x + y - (at_1^3 + 2at_1) = 0$

Putting $x = at^2, y = 2at$,

$at_1t^2 + 2at - (at_1^3 + 2at_1) = 0$

As t_1 is a real root of the equation, it has another real root

t_2 given by $t_1 + t_2 = \frac{-2a}{at_1}$

i.e. $t_2 = -\frac{2}{t_1} - t_1$

Further, $t_1t_2 = -(2 + t_1^2) < 0$

$\therefore t_2 \neq t_1$

(c) (i) Let $P_n = (at_n^2, 2at_n)$, where $t_n \neq 0$.

By (b), $x_{n+1} - x_n = at_n^2 - at_{n+1}^2 = a(t_n^2 - t_{n+1}^2)$
 $= a(t_n - t_{n+1})(t_n + t_{n+1})$
 $= a(t_n - t_{n+1}) \left(\frac{2+t_n^2}{t_n} \right)$
 $= \frac{4a}{t_n} - 4a$
 $= \frac{4a}{x_n} - 4a$

(ii) $x_{n+1} - x_n = \sum_{k=1}^n (x_{k+1} - x_k)$
 $= \sum_{k=1}^n \left(\frac{4a^2}{x_k} + 4a \right)$
 $> 4na$ as $x_k > 0$

As $n \rightarrow \infty, x_{n+1} \rightarrow \infty$ and therefore $\lim_{n \rightarrow \infty} \frac{1}{x_n} = 0$

(iii) $\lim_{n \rightarrow \infty} (x_{n+1} - x_n) = \lim_{n \rightarrow \infty} \left(\frac{4a^2}{x_n} + 4a \right) = 4a$

$|y_{n+1}| - |y_n| = 2\sqrt{a} (\sqrt{x_{n+1}} - \sqrt{x_n})$
 $= \frac{2\sqrt{a} (x_{n+1} - x_n)}{\sqrt{x_{n+1}} + \sqrt{x_n}}$

As $n \rightarrow \infty, x_n \rightarrow \infty$ and $\lim_{n \rightarrow \infty} (x_{n+1} - x_n) = 4a$,

$\lim_{n \rightarrow \infty} (|y_{n+1}| - |y_n|) = 0$

If cond. show $\Delta > 0$.
 avoid this mark in (c) when used

3(a) $a_{n+1} - a_n = \frac{r}{(n+1)!} - \frac{r}{n!} > 0$
 $\{a_n\}$ is monotonic increasing.

II

1

For $r \geq 2, r! \geq r(r-1)$

$\frac{1}{r!} \leq \frac{1}{r-1} - \frac{1}{r}$

$\sum_{r=0}^n \frac{1}{r!} = 2 + \sum_{r=2}^n \left(\frac{1}{r-1} - \frac{1}{r} \right)$

$= 3 - \frac{1}{n} < 3$

Hence $\{a_n\}$ is bounded above by 3 and is therefore convergent.

1

1

3

1

(b) $b_n = \sum_{r=0}^n C_r^n \frac{1}{n^r}$

$= 2 + \sum_{r=2}^n \frac{n(n-1)\dots(n-r+1)}{r! n^r}$

$= 2 + \sum_{r=2}^n \frac{1}{r!} \left(1 - \frac{1}{n}\right) \dots \left(1 - \frac{r-1}{n}\right)$

$\leq 2 + \sum_{r=2}^n \frac{1}{r!}$

$= a_n$

1

$\frac{1}{3}$

(c) Let n be fixed and greater than 1. We shall prove the inequality

by induction on r . The inequality holds for $r = 2$ as

$1 - \frac{(2)(1)}{n} \leq 1 - \frac{1}{n}$

Assume that it holds for some r where $2 \leq r < n$, then

$\left(1 - \frac{1}{n}\right) \dots \left(1 - \frac{r-1}{n}\right) \left(1 - \frac{r}{n}\right) \geq \left[1 - \frac{r(r-1)}{n}\right] \left(1 - \frac{r}{n}\right)$ ($1 - \frac{r}{n} \geq 0$)

$= 1 - \frac{r(r-1)}{n} - \frac{r}{n} + \frac{r^2(r-1)}{n^2}$

$> 1 - \frac{r^2}{n}$

$> 1 - \frac{r(r+1)}{n}$

the inequality holds for r where $2 \leq r \leq n$.

1

1

2

5 (a) (i) $\int_0^1 [f(x)]^2 dx = \int_0^1 \sum_{i=0}^n a_i x^i f(x) dx$

$$\sum_{i=0}^n a_i \int_0^1 x^i f(x) dx = a_0 \int_0^1 f(x) dx$$

as $\int_0^1 x^i f(x) dx = 0$ for $i = 1, 2, \dots, n$

(ii) For $k = 1, 2, \dots, n$

$$\int_0^1 x^k f(x) dx = 0$$

$$\int_0^1 x^k (a_n x^n + a_{n-1} x^{n-1} + \dots + a_0) dx = 0$$

$$\left[\frac{a_n}{k+n+1} x^{k+n+1} + \frac{a_{n-1}}{k+n} x^{k+n} + \dots + \frac{a_0}{k+1} x^{k+1} \right]_0^1 = 0$$

$$\frac{a_n}{k+n+1} + \frac{a_{n-1}}{k+n} + \dots + \frac{a_0}{k+1} = 0$$

(b) Let $\frac{a_n}{k+n+1} + \frac{a_{n-1}}{k+n} + \dots + \frac{a_0}{k+1} = \frac{Q(t)}{(t+n+1)(t+n)\dots(t+1)}$

where $Q(t)$ is a polynomial in t of degree $\leq n$.

By (a)(ii), $Q(t) = 0$ for $t = 1, 2, \dots, n$.

Hence $Q(t) = C(t-1)(t-2)\dots(t-n)$ for some constant C .

(c) Putting $t = 0$ in (b),

$$\frac{a_n}{n+1} + \frac{a_{n-1}}{n} + \dots + \frac{a_0}{1} = (-1)^n \frac{n! C}{(n+1)!}$$

But $\int_0^1 f(x) dx = \left[\frac{a_n}{n+1} x^{n+1} + \frac{a_{n-1}}{n} x^n + \dots + a_0 x \right]_0^1$

$$= \frac{a_n}{n+1} + \frac{a_{n-1}}{n} + \dots + a_0$$

The answers follows.

Multiplying both sides of (b) by $(t+1)$ and putting $t = -1$.

we have $a_0 = \frac{(-1)^n (n+1)! C}{n!} = (-1)^n (n+1)C$.

By (a), $\int_0^1 [f(x)]^2 dx = a_0 \int_0^1 f(x) dx$

$$= (-1)^n (n+1)C \int_0^1 f(x) dx = (n+1)C \int_0^1 f(x) dx$$

Solution

87 Marks Remarks

6(a) (i) Suppose l_1 and l_2 intersect at $\begin{pmatrix} \alpha \\ \beta \end{pmatrix} \exists c_1, c_2 \in \mathbb{R}$

such that $\begin{pmatrix} a_1 \\ b_1 \\ c_1 \end{pmatrix} + c_1 \begin{pmatrix} p_1 \\ q_1 \\ r_1 \end{pmatrix} = \begin{pmatrix} \alpha \\ \beta \\ 1 \end{pmatrix}$

$$= \begin{pmatrix} a_2 \\ b_2 \\ c_2 \end{pmatrix} + c_2 \begin{pmatrix} p_2 \\ q_2 \\ r_2 \end{pmatrix}$$

$$\begin{pmatrix} a_1 - a_2 \\ b_1 - b_2 \\ c_1 - c_2 \end{pmatrix} + c_1 \begin{pmatrix} p_1 \\ q_1 \\ r_1 \end{pmatrix} - c_2 \begin{pmatrix} p_2 \\ q_2 \\ r_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

The system $\begin{pmatrix} a_1 - a_2 & p_1 & p_2 \\ b_1 - b_2 & q_1 & q_2 \\ c_1 - c_2 & r_1 & r_2 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$ has a non-trivial

solution $\begin{pmatrix} 1 \\ c_1 \\ -c_2 \end{pmatrix}$

$$\begin{vmatrix} a_1 - a_2 & p_1 & p_2 \\ b_1 - b_2 & q_1 & q_2 \\ c_1 - c_2 & r_1 & r_2 \end{vmatrix} = 0$$

(ii) The vector $\begin{pmatrix} p_1 \\ q_1 \\ r_1 \end{pmatrix} \times \begin{pmatrix} p_2 \\ q_2 \\ r_2 \end{pmatrix}$ is normal to the plane containing

l_1 and l_2 .

$\therefore l_1$ and l_2 are distinct and they intersect, $\begin{pmatrix} p_1 \\ q_1 \\ r_1 \end{pmatrix} \times \begin{pmatrix} p_2 \\ q_2 \\ r_2 \end{pmatrix} \neq \mathbf{0}$

The plane containing l_1 and l_2 is given by

$$\begin{vmatrix} x - a_1 & y - b_1 & z - c_1 \\ p_1 & q_1 & r_1 \\ p_2 & q_2 & r_2 \end{vmatrix} = 0$$

$\vec{r} \cdot \vec{m} = 0$

$$\begin{vmatrix} x - a_1 & y - b_1 & z - c_1 \\ p_1 & q_1 & r_1 \\ p_2 & q_2 & r_2 \end{vmatrix} = 0$$

2

1

(b) (i) The required plane is parallel to the vector

$$\begin{pmatrix} q \\ r \\ p \end{pmatrix} \times \begin{pmatrix} r \\ p \\ q \end{pmatrix} = (rq - p^2)\underline{i} + (pr - q^2)\underline{j} + (pq - r^2)\underline{k}$$

As L_1, L_2, L_3 are concurrent at the origin, by (a)(ii), the equation of the plane is given by

$$\begin{vmatrix} x-0 & y-0 & z-0 \\ p & q & r \\ rq-p^2 & pr-q^2 & pq-r^2 \end{vmatrix} = 0$$

This can be written as

$$(pq + qr + rp)(q - r)x + (r - p)y + (p - q)z = 0$$

As p, q, r are distinct, this plane is uniquely defined if

$$pq + qr + rp \neq 0$$

Its equation is $(q - r)x + (r - p)y + (p - q)z = 0$.

(ii) If $pq + qr + rp = 0$, L_1, L_2, L_3 are mutually perpendicular.

There are infinitely many planes, each of which contains L_1 and which is perpendicular to the plane containing L_2 and L_3 .

The equation of one such plane is

$$\begin{vmatrix} x & y & z \\ p & q & r \\ q & r & p \end{vmatrix} = 0$$

7

$$\begin{aligned} \text{Let } f(x) &= \frac{x^2(x+7)}{x-1} = x^2 + 8x + 8 + \frac{8}{x-1} \\ f'(x) &= 2x + 8 - \frac{8}{(x-1)^2} = \frac{2x(x^2 + 2x - 7)}{(x-1)^2} \\ f''(x) &= 2 + \frac{16}{(x-1)^3} = \frac{2(x+1)(x^2 - 4x + 7)}{(x-1)^3} \end{aligned}$$

$$\text{As } f(x) = \begin{cases} g(x), & x > 0 \\ 0, & x = 0 \\ -g(x), & x < 0 \end{cases}$$

$$f'(x) = \begin{cases} g'(x), & x > 0 \\ 0, & x = 0 \\ -g'(x), & x < 0 \end{cases}$$

$$f''(x) = \begin{cases} g''(x), & x > 0 \\ -g''(x), & x < 0 \end{cases}$$

(b) (i) Putting $f'(x) = 0$, $x = 0$ or $-1 \pm 2\sqrt{2}$

At $x = -1 + 2\sqrt{2}$, $f''(x) > 0$
 $(-1 + 2\sqrt{2}, 13 + 16\sqrt{2})$ is a minimum point
for $(1.83, 35.63)$

At $x = -1 - 2\sqrt{2}$, $f''(x) < 0$
 $(-1 - 2\sqrt{2}, -13 + 16\sqrt{2})$ is a maximum point
for $(-3.83, 9.63)$

[The point $(0, 0)$ will be discussed in (ii)]

Next $x = 1$ is obviously an asymptote

As neither $\lim_{x \rightarrow \infty} \frac{f(x)}{x}$ nor $\lim_{x \rightarrow -\infty} \frac{f(x)}{x}$ exists, the graph of $f(x)$ has no oblique asymptotes.

(ii) Putting $f''(x) = 0$, $x = -1$

As $f''(x)$ changes sign at $x = -1$, $(-1, 3)$ is a point of inflexion.

Consider the point $(0, 0)$.

For $x \geq 0$ slightly, $f''(x) < 0$, $f'(x)$ is decreasing.


For $x < 0$ slightly, $f''(x) > 0$, $f'(x)$ is increasing.

$(0, 0)$ is another inflexion point.

As $f''(x)$ exists in \mathbb{R} except at $x = 0$ or 1 , these two are the only inflexion points.

1
2

7(c) The curve cuts the axes at (0, 0) and (-7, 0)



3/4

8(a) Integrating by parts,

$$\int_a^x (x-t)g''(t)dt = (x-t)g'(t) \Big|_a^x + \int_a^x g'(t)dt$$

$$= -(x-a)g'(a) + g(x) - g(a)$$

$$g(x) = g(a) + (x-a)g'(a) + \int_a^x (x-t)g''(t)dt \quad \forall x \in \mathbb{R}$$

(b) (i) If $g(x) = \int_a^x f(t)dt - \frac{(x-a)}{2}[f(x) + f(a)]$, g has a continuous

$$\text{second derivative and } g'(x) = f(x) - \frac{(x-a)}{2} f'(x) - \frac{1}{2}[f(x)+f(a)]$$

$$= -\frac{(x-a)}{2} f'(x) + \frac{1}{2}[f(x)-f(a)]$$

$$g''(x) = -\frac{(x-a)}{2} f''(x) \dots \dots \dots$$

$$\therefore \text{ by (a), } g(x) = 0 + (x-a)(0) + \int_a^x (x-t) \left[-\frac{(t-a)}{2} f''(t) \right] dt$$

$$= -\int_a^x \frac{(x-t)(t-a)}{2} f''(t) dt$$

$$(ii) \text{ Putting } x = b \text{ in (i) } |g(b)| = \left| \int_a^b \frac{(b-t)(t-a)}{2} f''(t) dt \right|$$

$$\leq \frac{M}{2} \left| \int_a^b (b-t)(t-a) dt \right|$$

$$= \frac{M}{2} \left| \left[-bt + \frac{(a+b)t}{2} - \frac{t^3}{3} \right]_a^b \right|$$

$$= \frac{M}{12} (b-a)^3$$

Dividing the interval $[0, 1]$ into n equal sub-intervals by the points $a_k = \frac{k}{n}, k = 0, 1, 2, \dots, n$, we have

$$\left| \int_{\frac{k}{n}}^{\frac{k+1}{n}} f(t) dt - \frac{1}{2} \cdot \frac{1}{n} \left[f\left(\frac{k+1}{n}\right) + f\left(\frac{k}{n}\right) \right] \right| \leq \frac{M}{12n^3}$$

$$\left| \int_0^1 f(t) dt - \frac{n-1}{2} \cdot \frac{1}{2n} \left[f\left(\frac{k+1}{n}\right) + f\left(\frac{k}{n}\right) \right] \right|$$

$$= \left| \sum_{k=0}^{n-1} \left[\int_{\frac{k}{n}}^{\frac{k+1}{n}} f(t) dt - \frac{1}{2n} \left[f\left(\frac{k+1}{n}\right) + f\left(\frac{k}{n}\right) \right] \right] \right|$$

$$\leq \sum_{k=0}^{n-1} \left| \int_{\frac{k}{n}}^{\frac{k+1}{n}} f(t) dt - \frac{1}{2n} \left[f\left(\frac{k+1}{n}\right) + f\left(\frac{k}{n}\right) \right] \right|$$

$$\leq \frac{M}{12n^2}$$

1/11

Solution

9(a) (i) Taking $y(x) = e^{\lambda x}$, $y'(x) = \lambda e^{\lambda x}$

$$h^{(n)}(x) = \frac{d^n}{dx^n} e^{(1+\lambda)x}$$

$$= (1+\lambda)^n e^{(1+\lambda)x}$$

$$f^{(k)}(x) e^{(n-k)x} = \left(\frac{d^k}{dx^k} e^{\lambda x} \right) \left(\frac{d^{n-k}}{dx^{n-k}} e^x \right)$$

$$= \lambda^k e^{\lambda x} \cdot e^x$$

$$= \lambda^k e^{(1+\lambda)x}, \quad k = 0, 1, \dots, n$$

$$(ii) (1+\lambda)^n e^{(1+\lambda)x} = \sum_{k=0}^n a_k \lambda^k e^{(1+\lambda)x}$$

$$(1+\lambda)^n = \sum_{k=0}^n a_k \lambda^k$$

Since this is true for any λ ,

$$a_k = \binom{n}{k} \lambda^k, \quad k = 0, 1, \dots, n$$

(b) (i) Using the above result,

$$y^{(n)}(x) = \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{m!}{k!} x^{n-k} e^{-x}$$

$$= \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{k!} x^{n-k} e^{-x}$$

$$= \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{k!} x^k e^{-x}$$

$$y(x) = \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{k!} x^k e^{-x}$$

which is a polynomial of degree n .

The coefficient of the term x^k is $\binom{n}{k} \frac{(-1)^k}{k!}$.

87 Marks Remarks

Solution

9(b) (ii) $xu'(x) + (x-m)u(x) = x[mx^{m-1}e^{-x} - x^m e^{-x}] + (x-m)x^m e^{-x} = 0$

Differentiating $(m+1)$ times,

$$\{xu'(x)\}^{(m+1)} = xu^{(m+2)}(x) + (m+1)u^{(m+1)}(x)$$

$$\{(x-m)u(x)\}^{(m+1)} = (x-m)u^{(m+1)}(x) + (m+1)u^{(m)}(x)$$

$$\therefore \text{L.S.} = xu^{(m+2)}(x) + (x+1)u^{(m+1)}(x) + (m+1)u^{(m)}(x)$$

$$= 0$$

$$(iii) y'(x) = e^x [u^{(m+1)}(x) + u^{(m)}(x)]$$

$$y''(x) = e^x [u^{(m+2)}(x) + 2u^{(m+1)}(x) + u^{(m)}(x)]$$

$$\therefore xy''(x) + (1-x)y'(x) + my(x)$$

$$= xe^x [u^{(m+2)}(x) + 2u^{(m+1)}(x) + u^{(m)}(x)] +$$

$$(1-x)e^x [u^{(m+1)}(x) + u^{(m)}(x)] + me^x u^{(m)}(x)$$

$$= e^x [xu^{(m+2)}(x) + (x+1)u^{(m+1)}(x) + (m+1)u^{(m)}(x)]$$

$$= 0 \quad \text{by (ii)}$$

87 Marks Remarks

1

1

1

for either

1

1

for either

1

1/10