

香港考試局

HONG KONG EXAMINATIONS AUTHORITY

一九八八年香港高級程度會考  
HONG KONG ADVANCED LEVEL EXAMINATION, 1988

PURE MATHEMATICS (PAPER 1)

MARKING SCHEME

This is a restricted document.

It is meant for use by markers of this paper for marking purposes only.

Reproduction in any form is strictly prohibited.

Special Note for Teacher Markers

It is highly undesirable that this marking scheme should fall into the hands of students. They are likely to regard it as a set of model answers, which it certainly is not.

Markers should therefore resist pleas from their students to have access to this document. Making it available would constitute misconduct on the part of the marker and is, moreover, in breach of the 1977 Hong Kong Examinations Authority Ordinance.

Solutions

88 Marks

Since  $\alpha$  is a root of  $P(x) - x = 0$ ,  $P(\alpha) - \alpha = 0$

i.e.  $P(\alpha) = \alpha$

$$P(P(\alpha)) - \alpha = P(\alpha) - \alpha$$

$$= 0$$

$\therefore \alpha$  is also a root of  $P(P(x)) - x = 0$ .

$$\begin{aligned} \text{(b)(i) } P(P(x)) - x &= (x^2 + ax + b)^2 + a(x^2 + ax + b) + b - x \\ &= x^4 + 2ax^3 + (a^2 + 2b + a)x^2 + (2ab + a^2 - 1)x + b^2 + ab + b \end{aligned}$$

By (a),  $x^2 + (a-1)x + b$  must be a factor of  $P(P(x)) - x$

By inspection,  $P(P(x)) - x = (x^2 + (a-1)x + b)(x^2 + (a+1)x + a + b + 1)$

(ii)  $P(P(x)) - x = 0$  has four real roots

$$\text{iff } (a-1)^2 \geq 4b$$

$$\text{and } (a+1)^2 \geq 4(a+b+1).$$

The second inequality can be written as  $(a-1)^2 \geq 4b + 4$

$$> 4b$$

$\therefore$  the required condition is  $(a-1)^2 \geq 4b + 4$ .

(c) Set  $P(x) = x^2 - 3x + 1$ . Two of the roots of  $P(x) - x = 0$ ,

and hence of  $P(P(x)) - x = 0$ , are  $\frac{4 \pm \sqrt{16-4}}{2}$  i.e.  $2 \pm \sqrt{3}$

The other two roots are given by  $x^2 + (-3+1)x + (-3+1+1) = 0$

$$x = 1 \pm \sqrt{2}$$

Alternative Solution

$$\begin{aligned} \text{(c) } (x^2 - 3x + 1)^2 - 3(x^2 - 3x + 1) + (1 - x) \\ = x^4 - 6x^3 + 6x^2 + 2x - 1 \end{aligned}$$

$$= (x^2 - 4x + 1)(x^2 - 2x - 1)$$

$$\text{The solutions are } x = \frac{4 \pm \sqrt{16-4}}{2}$$

$$\text{or } x = \frac{2 \pm \sqrt{4+4}}{2}$$

$$\text{i.e. } x = 2 \pm \sqrt{3} \text{ or } 1 \pm \sqrt{2}$$

1	
1	
2	
1	
2	
1	
1	
1	
1	
5	
1	
1	
1	
1	
4	

1+1  
4

88 Marks

2(a)  $Mx = \lambda x$  has a non-zero solution.

iff  $(M - \lambda I)x$  has a non-zero solution

iff  $|M - \lambda I| = 0$

Since  $|M - \lambda I| = 0$  is a cubic equation in  $\lambda$ , it has at most 3 solutions. i.e.  $M$  has at most 3 eigenvalues.

(b) Suppose  $\lambda$  is an eigenvalue of  $M$ .  $Mx = \lambda x$  for some non-zero  $x$ .

Then  $a_0 Ix = a_0 x$

$a_1 Mx = a_1 \lambda x$

$a_2 M^2 x = (a_2 M) \lambda x = a_2 \lambda^2 x$

etc.

$a_n M^n x = a_n \lambda^n x$

$(a_0 I + a_1 M + \dots + a_n M^n)x = a_0 Ix + a_1 Mx + \dots + a_n M^n x$   
 $= a_0 x + a_1 \lambda x + \dots + a_n \lambda^n x$   
 $= (a_0 + a_1 \lambda + \dots + a_n \lambda^n)x$

(c) (i)  $A = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$ ,  $|A - \lambda I| = 0$

iff  $\begin{vmatrix} -\lambda & 1 & 0 \\ 1 & -\lambda & 1 \\ 0 & 1 & -\lambda \end{vmatrix} = 0$

iff  $\lambda = 0$  or  $\pm\sqrt{2}$

By (a), these are all the eigenvalues of  $A$ .

(ii)  $A^2 = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 1 \end{pmatrix}$

Consider  $a_0 I + a_1 A + a_2 A^2$

$= \begin{pmatrix} a_2 + a_0 & a_1 & a_2 \\ a_1 & 2a_2 + a_0 & a_1 \\ a_2 & a_1 & a_2 + a_0 \end{pmatrix}$   
 $= B$

By (b), if  $\lambda$  is an eigenvalue of  $A$ , then  $a_0 + a_1 \lambda + a_2 \lambda^2$  is an eigenvalue of  $B$ .

Putting  $\lambda = 0, \pm\sqrt{2}$ , three eigenvalues of  $B$  are  $a_0, a_0 + a_1 \sqrt{2} + 2a_2,$

$a_0 - a_1 \sqrt{2} + 2a_2$

As  $a_0, a_1, a_2$  are non-zero integers, these eigenvalues are distinct and the answer follows from (a).

1  
7

3 (a) We shall prove by induction.

For  $n = 1$ ,  $p_2 q_1 - p_1 q_2 = (a_2 a_1 + 1) - a_1 a_2 = 1$   
 $= (-1)^2$

Assuming that  $p_{k+1} q_k - p_k q_{k+1} = (-1)^{k+1}$  for some  $k \geq 1$ ,

then  $p_{k+2} q_{k+1} - p_{k+1} q_{k+2} = (a_{k+2} p_{k+1} + p_k) q_{k+1} - p_{k+1} (a_{k+2} q_{k+1} + q_k)$   
 $= p_k q_{k+1} - p_{k+1} q_k$   
 $= (-1)^{k+2}$  by the induction hypothesis.

Hence the equality holds for all positive  $n$ .

(b) (i) For any  $n \geq 1$ ,  $b_{n+2} - b_n = \frac{p_{n+2}}{q_{n+2}} - \frac{p_n}{q_n}$

$= \frac{(a_{n+2} p_{n+1} + p_n) q_n - (a_{n+2} q_{n+1} + q_n) p_n}{q_{n+2} q_n} = \frac{a_{n+2} (p_{n+1} q_n - p_n q_{n+1})}{q_{n+2} q_n}$   
 $= \frac{(-1)^{n+1} a_{n+2}}{q_{n+2} q_n}$

As  $a_{n+2}, q_{n+2}, q_n$  are positive,  $b_{n+2} > b_n$  if  $n$  is odd  
 $b_{n+2} < b_n$  if  $n$  is even

Hence  $\{b_{2n-1}\}$  is strictly increasing and  $\{b_{2n}\}$  strictly decreasing.

(ii) For any  $n \geq 1$ ,  $b_{2n} - b_{2n-1} = \frac{p_{2n}}{q_{2n}} - \frac{p_{2n-1}}{q_{2n-1}}$   
 $= \frac{p_{2n} q_{2n-1} - p_{2n-1} q_{2n}}{q_{2n} q_{2n-1}}$   
 $= \frac{(-1)^{2n}}{q_{2n} q_{2n-1}} > 0$

If  $n$  is odd,  $b_{2n} > b_{2n-1}$  by (i)

If  $n$  is even,  $b_{2n} < b_{2n-1}$

(iii) As  $\{b_{2n-1}\}$  is strictly increasing and bounded above by  $b_2$ ,  $\{b_{2n}\}$  is strictly decreasing and bounded below by  $b_1$ , both sequences converge.

$\lim_{n \rightarrow \infty} (b_{2n} - b_{2n-1}) = \lim_{n \rightarrow \infty} \frac{(-1)^{2n}}{q_{2n} q_{2n-1}} = 0$

as  $\{q_n\}$  is a sequence of strictly increasing integers.

Thus  $\lim_{n \rightarrow \infty} b_{2n} = \lim_{n \rightarrow \infty} b_{2n-1}$

I

3

1

1

1

1

1

1

1+1

1

1

1

1  
11

) 2 for 1st  
) given  
) part



6. (a) (i) For any  $a, b \in \mathbb{R}$  and  $\underline{x}, \underline{y} \in \mathbb{R}^3$ ,

$$\begin{aligned} \phi_u(ax + by) &= \underline{u} \cdot (a\underline{x} + b\underline{y}) \\ &= a(\underline{u} \cdot \underline{x}) + b(\underline{u} \cdot \underline{y}) \\ &= a \phi_u(\underline{x}) + b \phi_u(\underline{y}) \end{aligned}$$

$\therefore \phi_u$  is linear.

(ii) If  $\phi_u(\underline{x}) = \underline{u} \cdot \underline{x} = 0$  for any  $\underline{x} \in \mathbb{R}^3$ , then  $\underline{u} \cdot \underline{u} = 0$   
 $\underline{u} = \underline{0}$

(iii) Given  $\underline{u}, \underline{v} \in \mathbb{R}^3$ , if  $\phi_u = \phi_v$ ,

then  $\underline{u} \cdot \underline{x} = \underline{v} \cdot \underline{x} \quad \forall \underline{x} \in \mathbb{R}^3$

$$\underline{u} \cdot \underline{x} - \underline{v} \cdot \underline{x} = 0$$

$$(\underline{u} - \underline{v}) \cdot \underline{x} = 0 \quad \forall \underline{x} \in \mathbb{R}^3$$

By (ii),

$$\underline{u} - \underline{v} = \underline{0} \quad \text{or} \quad \underline{u} = \underline{v}$$

(b) Given  $f: \mathbb{R}^3 \rightarrow \mathbb{R}$ ,

$$\text{let } \underline{n} = (f(\underline{i}), f(\underline{j}), f(\underline{k}))$$

Then for any  $\underline{x} = (x_1, x_2, x_3) \in \mathbb{R}^3$

$$\begin{aligned} \underline{n} \cdot \underline{x} &= f(\underline{i})x_1 + f(\underline{j})x_2 + f(\underline{k})x_3 \\ &= f(x_1\underline{i} + x_2\underline{j} + x_3\underline{k}) \quad \text{as } f \text{ is linear} \\ &= f(\underline{x}) \end{aligned}$$

If  $\exists \underline{n}' \in \mathbb{R}^3$  such that  $f(\underline{x}) = \underline{n}' \cdot \underline{x} \quad \forall \underline{x} \in \mathbb{R}^3$ ,

by (a)(iii),  $\underline{n}' = \underline{n}$ .

Hence the uniqueness

88 Marks Remarks

1

1

1

1

1

1

1

7

2

1

1

1

2

7

7. (a) (i)  $p(1) = \frac{1}{6}$

$$p(2) = \frac{1}{6} + \frac{1}{6} \times \frac{1}{6} \quad (= \frac{7}{36})$$

$$p(3) = \frac{1}{6} + \frac{1}{6} \times \frac{1}{6} \times 2 + \frac{1}{6} \times \frac{1}{6} \times \frac{1}{6} \quad (= \frac{49}{216})$$

$$(ii) p(4) = \frac{1}{6} + \frac{1}{6} (p(1) + p(2) + p(3))$$

$$(iii) \text{For } n > 6, p(n) = \frac{1}{6} (p(n-6) + p(n-5) + \dots + p(n-1))$$

(b) (i) For  $k < 0, p(k) = 0; p(0) = 1$ .

$$\text{For } k > 0, p(k) = \frac{1}{6} [p(k-6) + p(k-5) + \dots + p(k-1)]$$

$$\therefore \sum_{k=1}^n p(k) = \sum_{k=1}^n \frac{1}{6} [p(k-6) + p(k-5) + \dots + p(k-1)]$$

$$= \frac{1}{6} \left[ \sum_{k=1}^n p(k-6) + \sum_{k=1}^n p(k-5) + \dots + \sum_{k=1}^n p(k-1) \right]$$

$$= \frac{1}{6} \left[ \sum_{k=0}^{n-6} p(k) + \sum_{k=0}^{n-5} p(k) + \dots + \sum_{k=0}^{n-1} p(k) \right]$$

$$= \frac{1}{6} [ 6 \times \sum_{k=0}^n p(k) - 6p(n) - 5p(n-1) - 4p(n-2) - 3p(n-3) - 2p(n-4) - p(n-5) ]$$

$$\therefore p(n) + \frac{5}{6} p(n-1) + \frac{4}{6} p(n-2) + \frac{3}{6} p(n-3) + \frac{2}{6} p(n-4) + \frac{1}{6} p(n-5)$$

$$= \sum_{k=0}^n p(k) - \sum_{k=0}^n p(k) = 1$$

(ii) Since  $\lim_{n \rightarrow \infty} p(n)$  exists,

$$\left[ 1 + \frac{5}{6} + \frac{4}{6} + \frac{3}{6} + \frac{2}{6} + \frac{1}{6} \right] \lim_{n \rightarrow \infty} p(n) = 1$$

$$\therefore \lim_{n \rightarrow \infty} p(n) = \frac{2}{9}$$

88 Marks Remarks

1

1

1

2

5

1

2

2

1

1

1

1

9

May use induction

8. (a) (i) As  $\sum_{j=1}^5 \alpha_j = 1$ , .....

$$\alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5 \bar{b}_4$$

$$= -\alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5 (\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5)$$

$$= -(\alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5 + \alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5 + \alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5 + \alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5 + \alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5)$$

$$= -b_1$$

$$\alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5 \bar{b}_2$$

$$= -\alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5 (\alpha_1 \alpha_2 \alpha_3 + \alpha_1 \alpha_2 \alpha_4 + \alpha_1 \alpha_2 \alpha_5 + \alpha_1 \alpha_3 \alpha_4 + \alpha_1 \alpha_3 \alpha_5 + \alpha_1 \alpha_4 \alpha_5 + \alpha_2 \alpha_3 \alpha_4 + \alpha_2 \alpha_3 \alpha_5 + \alpha_2 \alpha_4 \alpha_5 + \alpha_3 \alpha_4 \alpha_5)$$

$$= -(\alpha_1 \alpha_4 \alpha_5 + \alpha_3 \alpha_4 \alpha_5 + \alpha_3 \alpha_4 \alpha_5 + \alpha_2 \alpha_5 + \alpha_2 \alpha_4 + \alpha_2 \alpha_3 + \alpha_1 \alpha_5 + \alpha_1 \alpha_4 + \alpha_1 \alpha_3 + \alpha_1 \alpha_2)$$

$$= -b_3$$

(ii)  $\sum_{j=1}^5 \alpha_j = 0 \Rightarrow b_4 = 0$

$\Rightarrow b_1 = 0$  by (i) as  $\alpha_j \neq 0$ .

Next  $0 = (\sum_{j=1}^5 \alpha_j)^2$

$$= \sum_{j=1}^5 \alpha_j^2 + 2b_3$$

$$\sum_{j=1}^5 \alpha_j^2 = 0 \Rightarrow b_3 = 0$$

$$\Rightarrow b_2 = 0$$

As  $b_5 = 1$  and  $b_0 = -\alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5$ , the result follows.

1

May be awarded below when used.

1

1

1

1

1

$\frac{1}{7}$

8 (b) Consider the fifth roots of the complex number  $\alpha = \alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5$ , which form a regular pentagon. ( $|\alpha| = 1$ ).

These are the roots of the equation  $z^5 - \alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5 = 0$ .

If  $\sum_{j=1}^5 \alpha_j = 0$ , by (a)(i),

$$(z - \alpha_1)(z - \alpha_2)(z - \alpha_3)(z - \alpha_4)(z - \alpha_5) = z^5 - \alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5 = 0$$

iff  $z = \alpha_1, \alpha_2, \alpha_3, \alpha_4$  or  $\alpha_5$

$\therefore$  the fifth roots are exactly the  $\alpha_j$ 's.

On the other hand, if the  $\alpha_j$ 's form a regular pentagon,

without loss of generality, let  $\alpha_1 = \cos \theta + i \sin \theta$ ,

$$\alpha_2 = \cos(\theta + \frac{2\pi}{5}) + i \sin(\theta + \frac{2\pi}{5}) = \alpha_1 w$$

$$\alpha_3 = \alpha_1 w^2, \alpha_4 = \alpha_1 w^3 \text{ and } \alpha_5 = \alpha_1 w^4, \text{ where } w = \cos \frac{2\pi}{5} + i \sin \frac{2\pi}{5}$$

$$\alpha_1 + \alpha_2 + \dots + \alpha_5 = \alpha_1(1 + w + w^2 + w^3 + w^4)$$

Now  $w^5 = 1 \Rightarrow w^5 - 1 = 0$

$$\Rightarrow (w - 1)(1 + w + w^2 + w^3 + w^4) = 0$$

$$\Rightarrow 1 + w + w^2 + w^3 + w^4 = 0 \text{ as } w \neq 1.$$

$$\sum_{j=1}^5 \alpha_j = 0$$

Similarly, writing  $\alpha_j^2 = \alpha_1^2 w^{2(j-1)}$ ,  $j = 1, 2, \dots, 5$ .

$$\sum_{j=1}^5 \alpha_j^2 = 0, \text{ noting that } 1 + w^2 + w^4 + w^6 + w^8 = 0.$$

Alternative solution to the 'only if' part  
If the  $\alpha_j$ 's form a regular pentagon, they are the fifth roots of some complex number

$\alpha = a + bi$ ,  $a, b \in \mathbb{R}$ , i.e. the roots of the equation  $z^5 - \alpha = 0$ .  
Let  $z = \cos \theta + i \sin \theta$  (as  $|z| = 1$ ).

$$z^5 - \alpha = 0 \text{ iff } (\cos \theta + i \sin \theta)^5 - (a + bi) = 0$$

$$\text{iff } (\cos^5 \theta - 10 \cos^3 \theta \sin^2 \theta + 5 \cos \theta \sin^4 \theta - a) + (5 \cos^4 \theta \sin \theta - 10 \cos^2 \theta \sin^3 \theta + \sin^5 \theta - b)i = 0$$

$$\text{iff } (16 \cos^5 \theta - 20 \cos^3 \theta + 5 \cos \theta - a) + (16 \sin^5 \theta - 20 \sin^3 \theta + 5 \sin \theta - b)i = 0$$

As the coefficient of the term  $\cos \theta$  in the equation  $16 \cos^5 \theta - 20 \cos^3 \theta + 5 \cos \theta - a = 0$

is zero, the sum of the roots  $\cos \theta_1 + \cos \theta_2 + \dots + \cos \theta_5 = 0$ . Similar consideration

of the imaginary part gives  $\sin \theta_1 + \sin \theta_2 + \dots + \sin \theta_5 = 0$ .

$\therefore$  the sum of the roots of the equation  $z^5 - \alpha = 0$  is zero.

Next if  $\alpha_j$ 's form a regular pentagon,  $\alpha_j^2$ 's also form a regular pentagon

and  $\sum_{j=1}^5 \alpha_j^2 = 0$

1

1

1

1

1

1

1

$\frac{1}{7}$

1

1

1

1

1










(a)  $\angle PBD = \theta$

$BP = a\theta$   
 $x = OD + PE$   
 $= a(\cos\theta + \theta\sin\theta)$   
 $y = BD - BE$   
 $= a(\sin\theta - \theta\cos\theta), \theta \geq 0.$



(b)  $\frac{dy}{d\theta} = a\theta\sin\theta, \frac{dx}{d\theta} = a\theta\cos\theta$   
 $\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta}$   
 $= \tan\theta$

The slope of the tangent line at any point P on the locus

$= \tan\theta$   
 $= \text{slope of } OB.$

As  $PB \perp OB$ , the thread is normal to the locus

(c) The area bounded  $= \int_0^a x dy$   
 $= \int_0^{\pi/2} a^2(\cos\theta + \theta\sin\theta)\theta\sin\theta d\theta$   
 $= a^2 \int_0^{\pi/2} (\theta\sin\theta\cos\theta + \theta^2\sin^2\theta) d\theta$

Now  $\int_0^{\pi/2} \theta\sin\theta\cos\theta d\theta = \int_0^{\pi/2} \theta\sin 2\theta d\theta$   
 $= -\frac{1}{4} \int_0^{\pi/2} \theta d\cos 2\theta$   
 $= -\frac{1}{4} [\theta\cos 2\theta \Big|_0^{\pi/2} - \int_0^{\pi/2} \cos 2\theta d\theta]$   
 $= \frac{\pi^2}{16}$

$\int_0^{\pi/2} \theta^2\sin^2\theta d\theta = \int_0^{\pi/2} \theta^2 \left( \frac{1 - \cos 2\theta}{2} \right) d\theta$   
 $= \int_0^{\pi/2} \frac{\theta^2}{2} d\theta - \frac{1}{4} \int_0^{\pi/2} \theta^2 \cos 2\theta d\theta$   
 $= \frac{\pi^3}{48} - \frac{1}{4} \left[ \theta^2\sin 2\theta \Big|_0^{\pi/2} - \int_0^{\pi/2} 2\theta\sin 2\theta d\theta \right]$   
 $= \frac{\pi^3}{48} + \frac{\pi}{8}$

the area bounded  $= \frac{\pi a^2(12 + \pi^2)}{48}$  ( $\approx 1.43a^2$ )

Marks

II

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

(a) Let  $f(x) = ax^2 + bx + c$   
 $f'(x) = 2ax + b$

L.S.  $= (x - t)(ax + at + b)$

R.S.  $= (x - t)[2a(\frac{x+t}{2}) + b] = R.S.$

(b) (i) Differentiating both sides with respect to x

$g'(x) = \frac{1}{2}(x - t)g''(\frac{x+t}{2}) + g'(\frac{x+t}{2})$

$g''(x) = \frac{1}{4}(x - t)g'''(\frac{x+t}{2}) + g''(\frac{x+t}{2})$

$\therefore g'''(\frac{x+t}{2}) = \frac{4(g''(x) - g''(\frac{x+t}{2}))}{x - t}, x \neq t.$

(ii) By L'Hospital's Rule

$\lim_{x \rightarrow t} g'''(\frac{x+t}{2}) = \lim_{x \rightarrow t} \frac{4[g''(x) - g''(\frac{x+t}{2})]}{x - t} = \frac{0}{0}$

By the continuity of  $g'''$ ,

$g'''(t) = \lim_{x \rightarrow t} \frac{4[g'''(x) - \frac{1}{2}g'''(\frac{x+t}{2})]}{1}$

$= 2g'''(t)$

$\therefore g'''(t) = 0 \quad \forall t \in \mathbb{R}.$

Since  $g'''(t) = 0$ ,  $g''(t) = c_1$

$g'(t) = c_1t + c_2$

$g(t) = \frac{c_1}{2}t^2 + c_2t + c_3$

Hence g is a polynomial of degree  $\leq 2$ .

6. (a) For  $x > 0$ ,  $f'(x) = \frac{2x(4x+3)}{3(x+1)^2}$ ,  $f''(x) = \frac{2(20x^2+30x+9)}{9(x+1)^3}$

For  $x < 0$ ,  $x \neq -1$ ,  $f'(x) = \frac{-2x(4x+3)}{3(x+1)^2}$ ,  $f''(x) = \frac{-2(20x^2+30x+9)}{9(x+1)^3}$


At  $x = 0$ ,  $f'(x)$  exists and equals zero  
 $f''(x)$  doesn't exist ( $f''(0) = 2$  but  $f''(0) = -2$ )  
 (As  $x \rightarrow -1^-$ ,  $f'(x) \rightarrow \infty$ )  
 At  $x = -1$ ,  $f'(x)$  doesn't exist  
 (As  $x \rightarrow -1^+$ ,  $f'(x) \rightarrow -\infty$ )

$f''(x)$  also doesn't exist there.

(b)  $f'(x) = 0$  iff  $x = 0$  or  $-\frac{3}{4}$   
 $f''(x) = 0$  iff  $x = \frac{-15 \pm 3\sqrt{5}}{20}$  (-0.41, -1.09)

Noting that  $f$  is continuous in  $\mathbb{R}$ , the following could be concluded:

$x$	$x_1 = \frac{-15 - 3\sqrt{5}}{20}$	-1	$-\frac{3}{4}$	$x_2 = \frac{-15 + 3\sqrt{5}}{20}$	0
$y'$	+	not defined	-	+	+
$y''$	0	not defined	+	0	not defined
$y$	Inf pt.	max	min	Inf pt.	Inf pt.



(c)

3

Such great details not expected

5

(a)  $\int_0^{2\pi} (\cos px \cos qx + \sin px \sin qx) dx = \int_0^{2\pi} \cos(p-q)x dx$   
 $= \begin{cases} 2\pi & \text{if } p = q \\ 0 & \text{if } p \neq q \end{cases}$

(b)  $\left| \sum_{p=0}^n a_p (\cos x + i \sin x)^p \right|^2 = \left| \sum_{p=0}^n a_p (\cos px + i \sin px) \right|^2$   
 $= \left( \sum_{p=0}^n a_p \cos px \right)^2 + \left( \sum_{p=0}^n a_p \sin px \right)^2$   
 $= \sum_{p=0}^n \sum_{q=0}^n a_p a_q (\cos px \cos qx + \sin px \sin qx)$   
 $= \sum_{p=0}^n \sum_{q=0}^n a_p a_q (\cos px \cos qx + \sin px \sin qx)$

$\int_0^{2\pi} |f(x)|^2 dx = \int_0^{2\pi} \left[ \sum_{p=0}^n \sum_{q=0}^n a_p a_q (\cos px \cos qx + \sin px \sin qx) \right] dx$   
 $= \sum_{p=0}^n \sum_{q=0}^n a_p a_q \int_0^{2\pi} (\cos px \cos qx + \sin px \sin qx) dx$   
 $= 2\pi \sum_{p=0}^n a_p^2$  by (a)

(c)  $(1 + \cos x + i \sin x)^n = [1 + (\cos x + i \sin x)]^n = \sum_{p=0}^n \binom{n}{p} (\cos x + i \sin x)^p$

By (b)  $\int_0^{2\pi} |g(x)|^2 dx = 2\pi \sum_{p=0}^n \binom{n}{p}^2$

Next,  $(1 - \cos x - i \sin x)^n = \sum_{p=0}^n \binom{n}{p} (-1)^p (\cos x + i \sin x)^p$

$\int_0^{2\pi} |h(x)|^2 dx = \int_0^{2\pi} |g(x)|^2 dx$

(d) The coefficient of  $x^n$  in the expansion of  $(1+x)^{2n} = \binom{2n}{n}$

On the other hand,  $(1+x)^{2n} = (1+x)^n (1+x)^n = \left( \sum_{p=0}^n \binom{n}{p} x^p \right) \left( \sum_{p=0}^n \binom{n}{p} x^p \right)$   
 $= \left( \sum_{p=0}^n \binom{n}{p} x^p \right) \left( \sum_{p=0}^n \binom{n}{p} x^p \right)$

The coefficient of the term  $x^n = \sum_{p=0}^n \binom{n}{p}^2$

Hence the result.

3

5

2

1

1

4

1

1

1

3

8. (a) Substitute  $x, y, z$  of  $L_2$  into  $L_1, 1 + 2t + 1 = 0 \Rightarrow t = -1$   
 $1 + 1 + t = 0 \Rightarrow t = -2$ ,  
 which are inconsistent.  $L_1$  and  $L_2$  therefore do not intersect each other.  
 Further, putting  $x = t$  in  $L_1, y = -t, z = t$ , which are the parametric equations of  $L_1$ .  
 Comparing the direction numbers of  $L_1$  and  $L_2$ , we see that they are not parallel.  
 Hence they are not coplanar.

(b) The system of planes containing  $L_1$  is given by  
 $x + y + \lambda(y + z) = 0$  or  $x + (1 + \lambda)y + \lambda z = 0$ .

For one of these planes to be parallel to  $L_2$ ,  
 $(1 + \lambda, \lambda) \cdot (2, 0, 1) = 0$   
 $\lambda = -2$

$\therefore \pi_1$  is given by  $x - y - 2z = 0$ .

(c) Let the equation of  $\pi_2$  be  $Ax + By + Cz + D = 0$ .

Since it contains  $L_2, (2, 0, 1) \cdot (A, B, C) = 0$   
 or  $2A + C = 0$ .

As  $\pi_1$  and  $\pi_2$  are perpendicular,  $(1, -1, -2) \cdot (A, B, C) = 0$   
 or  $A - B - 2C = 0$ .

Solving these two equations,  $A : B : C = 1 : 5 : -2$ .

Writing  $\pi_2$  as  $x + 5y - 2z + D = 0$  and putting  $(x, y, z) = (1, 1, 1)$ ,  
 which is a point on  $L_2, D = -4, \therefore \pi_2$  is given by  $x + 5y - 2z - 4 = 0$ .

(d) The vector  $(1, -1, -2)$  is perpendicular to  $\pi_1$ .

Let  $O = (1+t, 1-t, 1-2t)$

Substituting in  $\pi_1, (1+t) - (1-t) - 2(1-2t) = 0$   
 $t = \frac{1}{3}$

$\therefore O = (\frac{4}{3}, \frac{2}{3}, \frac{1}{3})$

The shortest distance between  $L_1$  and  $L_2$

$$= \sqrt{(1 - \frac{4}{3})^2 + (1 - \frac{2}{3})^2 + (1 - \frac{1}{3})^2}$$

$$= \sqrt{\frac{2}{3}}$$

II

$\frac{1}{3}$

$\frac{1}{3}$

$\frac{1}{4}$

1

1

(a) If  $f$  is Lipschitz-continuous,  $0 \leq |f(x_1) - f(x_2)| \leq k|x_1 - x_2|$  for any  $x_1, x_2 \in I$ .

Since  $\lim_{x_1 \rightarrow x_2} k|x_1 - x_2| = 0, \lim_{x_1 \rightarrow x_2} |f(x_1) - f(x_2)| = 0$ .

$\therefore \lim_{x_1 \rightarrow x_2} f(x_1) = f(x_2)$  and  $f$  is continuous at any  $x_2 \in I$ .

For any  $x_1, x_2 \in (0, 1), |g(x_1) - g(x_2)|$   
 $= |\sqrt{x_1} - \sqrt{x_2}|$   
 $= \frac{1}{\sqrt{x_1} + \sqrt{x_2}} |x_1 - x_2|$

If  $x_1$  and  $x_2$  tend to zero,  $\frac{1}{\sqrt{x_1} + \sqrt{x_2}}$  increases without bound. Hence the

Lipschitz condition cannot be satisfied.

(b) For any  $x_1, x_2 \in I$ , we may let  $x_1 \geq x_2$ . As  $f(x)$  is continuous,

$$|f(x_1) - f(x_2)| = \left| \int_{x_2}^{x_1} f'(t) dt \right|$$

$$\leq \int_{x_2}^{x_1} |f'(t)| dt$$

$$\leq \int_{x_2}^{x_1} M dt$$

$$= M|x_1 - x_2|$$

i.e.  $f$  is Lipschitz-continuous on  $I$ .

(c) (i) Consider the function  $h(x) = x - f(x)$ . Since  $f$  is continuous, by (a),

$h$  is also also continuous.

If  $f(a) = a$  or  $f(b) = b$ , we are through.

Otherwise, let  $a < f(a), f(b) < b$ . Then  $h(a) < 0$  and  $h(b) > 0$ .

$\therefore \exists x_0 \in (a, b)$  such that  $h(x_0) = 0$  i.e.  $x_0 - f(x_0) = 0$ .

(ii) Let  $x_0'$  be in  $(a, b)$  such that  $x_0' - f(x_0') = 0$ .

Since  $f$  is Lipschitz-continuous with  $0 < k < 1, k|x_0' - x_0| \geq |f(x_0') - f(x_0)|$   
 $= |x_0' - x_0|$

The inequality holds only if  $x_0' = x_0$ .  $\therefore$  the solution is unique.

1

1

1

1

4

1

1

1

3

1

1

1

1

1

1

1

1

May use mean value theorem